

ON THE CARDINALITY OF URYSOHN SPACES
AND WEAKLY H -CLOSED SPACES

FORTUNATA AURORA BASILE, Messina, NATHAN CARLSON, Thousand Oaks

Received March 21, 2018. Published online November 27, 2018.

Communicated by J. Gutiérrez García

Abstract. We introduce the cardinal invariant $\theta\text{-}aL'(X)$, related to $\theta\text{-}aL(X)$, and show that if X is Urysohn, then $|X| \leq 2^{\theta\text{-}aL'(X)\chi(X)}$. As $\theta\text{-}aL'(X) \leq aL(X)$, this represents an improvement of the Bella-Cammaroto inequality.

We also introduce the classes of firmly Urysohn spaces, related to Urysohn spaces, strongly semiregular spaces, related to semiregular spaces, and weakly H -closed spaces, related to H -closed spaces.

Keywords: Urysohn space; θ -closure; pseudocharacter; almost Lindelöf degree; cardinality; cardinal invariant

MSC 2010: 54A25, 54D10, 54D20

1. INTRODUCTION

We follow the notation from [8] and [9]. Recall that a *space* X is *Urysohn* if for every two distinct points $x, y \in X$ there are open sets U and V such that $x \in U$, $y \in V$ and $\overline{U} \cap \overline{V} = \emptyset$.

Many researchers have worked on the cardinality of Urysohn and H -closed spaces, in particular considering cardinal invariants defined by using the θ -closure operator defined below (see, for instance, [1], [2], [4], [5], [6], [7], [10], [11]).

For a space X , we denote by $\chi(X)$ ($\psi(X)$, $\pi\chi(X)$, $c(X)$, $t(X)$) the *character* (respectively, *pseudocharacter*, π -*character*, *cellularity*, *tightness*), see [8].

The θ -*closure* of a set A in a space X is the set $\text{cl}_\theta(A) = \{x \in X : \text{for every neighborhood } U \ni x, \overline{U} \cap A \neq \emptyset\}$; A is said to be θ -*closed* if $A = \text{cl}_\theta(A)$, see [13].

The authors wish to thank Jack Porter for his helpful comments and suggestions in several places in this paper.

The complement of a θ -closed set is called a θ -open set. Considering the fact that the θ -closure operator is not in general idempotent, Bella and Cammaroto defined in [4] the θ -closed hull of a subset A of a space X , denoted by $[A]_\theta$, that is the smallest θ -closed subset of X containing A .

The γ -closure (see [2]) of the set A in a space X is the set $\text{cl}_\gamma(A) = \{x: \text{for every open neighborhood } U \text{ of } x, \text{cl}_\theta(\overline{U}) \cap A \neq \emptyset\}$. A is said to be γ -closed if $A = \text{cl}_\gamma(A)$. The γ -closure operator is not in general idempotent.

If X is a Urysohn space, the θ -pseudocharacter of a point x in X (see [2]) is $\psi_\theta(x, X) = \min\{|\mathcal{U}|: \mathcal{U} \text{ is a family of open neighborhoods of } x \text{ and } \{x\} \text{ is the intersection of the } \theta\text{-closures of the closures of } \mathcal{U}\}$; the θ -pseudocharacter of X is $\psi_\theta(X) = \sup\{\psi_\theta(x, X): x \in X\}$.

The almost Lindelöf degree of a subset Y of a space X is $aL(Y, X) = \min\{k: \text{for every cover } \mathcal{V} \text{ of } Y \text{ consisting of open subsets of } X, \text{there exists } \mathcal{V}' \subseteq \mathcal{V} \text{ such that } |\mathcal{V}'| \leq k \text{ and } \bigcup\{\overline{V}: V \in \mathcal{V}'\} = Y\}$. The function $aL(X, X)$ is called the almost Lindelöf degree of X and denoted by $aL(X)$ (see [14] and [9]). The almost Lindelöf degree of X with respect to closed subsets is $aL_C(X) = \min\{k: \text{for every closed set } Y \subset X \text{ and every cover } \mathcal{V} \text{ of } Y \text{ consisting of open subsets of } X, \text{there exists } \mathcal{V}' \subseteq \mathcal{V} \text{ such that } |\mathcal{V}'| \leq k \text{ and } \bigcup\{\overline{V}: V \in \mathcal{V}'\} = Y\}$. The almost Lindelöf degree of X with respect to θ -closed subsets is $aL_\theta(X) = \min\{k: \text{for every } \theta\text{-closed set } Y \subset X \text{ and every cover } \mathcal{V} \text{ of } Y \text{ consisting of open subsets of } X, \text{there exists } \mathcal{V}' \subseteq \mathcal{V} \text{ such that } |\mathcal{V}'| \leq k \text{ and } \bigcup\{\overline{V}: V \in \mathcal{V}'\} = Y\}$. We have that the almost Lindelöf degree is hereditary with respect to θ -closed sets, so we have that $aL(X) = aL_\theta(X)$.

A space X is called H -closed if it has finite almost Lindelöf degree.

Let $(X, \tau) = X$ be a topological space. We say that a subset A of X is regular open (regular closed) if $A = \text{int}(\overline{A})$ ($A = \overline{\text{int}(A)}$). The family $\mathcal{B} = \{U: U \text{ is regular open in } X\}$ is a base for X . Space X equipped with the topology generated by the base \mathcal{B} is called the semiregularization of X and is denoted by X_s . If $X = X_s$, then X is called semiregular.

For a subset A of a space X we will denote by $[A]^{\leq \lambda}$ the family of all subsets of A of cardinality $\leq \lambda$.

In Section 2 we introduce the notions of weakly H -closed spaces, strongly semiregular spaces and firmly Urysohn spaces and we prove that if X is a H -closed, strongly semiregular, firmly Urysohn space, then X is compact.

In Section 3 we construct two new cardinal invariants denoted by $\theta\text{-}aL'(X)$, related to the θ -almost Lindelöf degree, and by $t_{\tilde{c}}(X)$, related to tightness, using an operator denoted by $\tilde{c}(\cdot)$ (for this new operator we have that if X is a space and $A \subset X$, then $\text{cl}_\theta(A) \subseteq \tilde{c}(A) \subseteq \text{cl}_\gamma(A)$). We prove some results concerning weakly H -closed spaces and, in particular, we prove that if X is a Urysohn space, then

$|X| \leq 2^{\theta-aL'(X)t_\varepsilon(X)\psi_\theta(X)}$, which represents an improvement of the Bella-Cammaroto inequality (see [4]).

2. WEAKLY H -CLOSED SPACES AND STRONG SEMIREGULAR SPACES

In [2] the cardinal invariant θ - $aL(X)$ was introduced, known as the θ -almost Lindelöf degree of the space X .

Definition 2.1. The θ -almost Lindelöf degree of a subset Y of a space X is θ - $aL(Y, X) = \min\{k: \text{for every cover } \mathcal{V} \text{ of } Y \text{ consisting of open subsets of } X, \text{ there exists } \mathcal{V}' \subseteq \mathcal{V} \text{ such that } |\mathcal{V}'| \leq k \text{ and } \bigcup\{\text{cl}_\theta(\overline{V}): V \in \mathcal{V}'\} = Y\}$.

The function θ - $aL(X, X)$ is the θ -almost Lindelöf degree of the space X and it is denoted by θ - $aL(X)$.

We have that θ - $aL(X) \leq aL(X)$ for every space X , and using a slight modification of Example 2.3 in [3] we prove that the previous inequality can be strict.

We say that a space is *weakly H -closed* if it has finite θ -almost Lindelöf degree. We obviously have that an H -closed space is weakly H -closed but the converse is not true as the following example shows:

Example 2.1. We consider Bing's tripod space \mathbb{B} . Let \mathbb{B} be the set $\{(x, y) \in \mathbb{Q}^2: y \geq 0\}$ with the topology generated by the neighborhood base $\mathcal{V}_{(x,y)} = \{N_\varepsilon(x, y): \varepsilon > 0\}$, where $N_\varepsilon(x, y) = \{(x, y)\} \cup (x - \frac{1}{\sqrt{3}}y - \varepsilon, x - \frac{1}{\sqrt{3}}y + \varepsilon) \times \{0\} \cup (x + \frac{1}{\sqrt{3}}y - \varepsilon, x + \frac{1}{\sqrt{3}}y + \varepsilon) \times \{0\}$. It is easily seen that X is not Urysohn and not H -closed but considering the fact that the θ -closure of the closure of every open set is the whole space we certainly have that X is weakly H -closed.

We also introduce a new cardinal function related to the weakly Lindelöf degree called the θ -weakly Lindelöf degree.

Definition 2.2. Let X be a topological space and Y a subset of X , the θ -weakly Lindelöf degree of Y with respect to X , denoted by θ - $wL(Y, X)$, is

$$\theta$$
- $wL(Y, X) = \min\left\{k: \text{for every open cover } \mathcal{U} \text{ of } Y \text{ by open subsets of } X \right. \\ \left. \text{there exists } \mathcal{V} \in [\mathcal{U}]^{\leq k} \text{ such that } Y = \text{cl}_\theta(\overline{\bigcup \mathcal{V}})\right\}.$

Definition 2.3. Let X be a topological space, the θ -weakly Lindelöf degree of X , denoted by θ - $wL(X)$, is θ - $wL(X, X)$.

Now we prove that for a θ -open subset U of a space X , θ - $wL(\overline{U}, X) \leq \theta$ - $wL(X)$.

Proposition 2.1. If U is a θ -open subset of a space X , then θ - $wL(\overline{U}, X) \leq \theta$ - $wL(X)$.

Proof. Let $k = \theta\text{-}wL(X)$, $U \subseteq X$ θ -open and \mathcal{U} a cover of \overline{U} by sets open in X . Then $\mathcal{U} \cup \{X \setminus \overline{U}\}$ is an open cover of X and since $k = \theta\text{-}wL(X)$ there exists $\mathcal{V} \in [\mathcal{U}]^{\leq k}$ such that $X = \text{cl}_\theta(X \setminus \overline{U}) \cup \text{cl}_\theta(\bigcup \mathcal{V})$. We want to prove that $U \cap \text{cl}_\theta(X \setminus \overline{U}) = \emptyset$. Suppose that there exists $x \in U \cap \text{cl}_\theta(X \setminus \overline{U})$. Point x is in U which is θ -open, so there exists an open set W such that $x \in W \subseteq \overline{W} \subseteq U$. Then $\overline{W} \cap X \setminus \overline{U} \neq \emptyset$, so $U \cap X \setminus \text{int}(\overline{U}) \neq \emptyset$, a contradiction. So U is contained in $\text{cl}_\theta(\bigcup \mathcal{V})$ and thus $\overline{U} \subseteq \text{cl}_\theta(\bigcup \mathcal{V})$, therefore $\theta\text{-}wL(\overline{U}, X) \leq k$. \square

Corollary 2.1. *If X is weakly H -closed and U is θ -open in X , then for every open cover \mathcal{U} of \overline{U} there exist a finite subfamily \mathcal{V} such that $\overline{U} \subseteq \bigcup_{V \in \mathcal{V}} \text{cl}_\theta(\overline{V})$.*

Definition 2.4. We say that a space X is *strongly semiregular* if the family $\mathcal{B} = \{X \setminus \overline{U} : U \text{ is a } \theta\text{-open subset of } X\}$ is a base for X .

It could be seen that \mathcal{B} satisfies the requirements to be a base for some topology. X equipped with the topology generated by the base \mathcal{B} is called the *strong semiregularization* of X and is denoted by X_{ss} .

It is natural to investigate the relation between the strong semiregularization X_{ss} and the semiregularization X_s of the space X .

Considering the fact that a θ -open set is open, the closure of an open set is regular closed and that the complement of a regular closed set is regular open, we have

Proposition 2.2. *A strongly semiregular space is semiregular.*

The converse of the above statement is not true.

Example 2.2. Let \mathbb{B}_s be the semiregularization of the space \mathbb{B} of Example 2.1. We have that if U is a θ -open subset of \mathbb{B} , then $\mathbb{B}_s \setminus U$ is nowhere dense and so it has empty interior, hence the space is not strongly semiregular.

We can also observe that for a Hausdorff space X , it may happen that X_{ss} is not Hausdorff even though X_s is Hausdorff.

Example 2.3. \mathbb{B}_{ss} has the indiscrete topology whereas \mathbb{B}_s is Hausdorff.

We now give the definition of a new axiom of separation.

Definition 2.5. A topological space X is *firmly Urysohn* if for every $x, y \in X$ with $x \neq y$, there exist two open subsets U, V of X such that $x \in U, y \in V$ and $\overline{U} \cap \text{cl}_\theta(\overline{V}) = \emptyset$.

We know that a firmly Urysohn space is Urysohn but we ask:

Question 2.1. Does there exist a Urysohn not firmly Urysohn space?

We investigate the θ -closed subsets of a firmly Urysohn, weakly H -closed space and we prove:

Proposition 2.3. *If X is firmly Urysohn and weakly H -closed and U is a θ -open subset of X , then $H = \overline{U}$ is θ -closed.*

Proof. Fix $x \notin H$. For each $y \in H$, there are open sets U_y and V_y such that $x \in U_y$, $y \in V_y$, and $\text{cl}_\theta(U_y) \cap \text{cl}_\theta(\overline{V_y}) = \emptyset$. By Corollary 2.1, for the open cover $\{V_y: y \in H\}$ of H , there is a finite subset $F \subseteq H$ such that $H \subseteq \bigcup_{y \in F} \text{cl}_\theta(\overline{V_y})$. Let $U = \bigcap_{y \in F} U_y$. Then $U \cap \bigcap_{y \in F} \text{cl}_\theta(\overline{V_y}) = \emptyset$. Thus, H is θ -closed. \square

We also investigate the relation between θ -open sets and regular spaces and we prove:

Proposition 2.4. *X has a base of θ -open sets if and only if X is regular.*

Proof. Suppose that X has a base consisting of θ -open sets. Let U be an open subset of X and $x \in U$, then there exists a θ -open subset V of X such that $x \in V \subseteq U$ and $x \notin X - V$. Subset V is θ -closed, so there exists an open subset W of X such that $x \in W$ and $\overline{W} \cap (X - V) = \emptyset$. This means $x \in W \subseteq \overline{W} \subseteq V \subseteq U$ if and only if X is a regular space. \square

Using the previous property, we find a connection with compact spaces. We recall an important result:

Lemma 2.1 ([12]). *A space X is H -closed and regular if and only if X is compact.*

If X is regular, then the θ -closure equals the closure and for this reason we can say that in a regular space the almost Lindelöf degree and the θ -almost Lindelöf degree coincide.

From Corollary 4.8(k) in [12], we prove:

Theorem 2.1. *If X is a firmly Urysohn, weakly H -closed and strongly semiregular space, then it is compact.*

Proof. Suppose that X is firmly Urysohn, weakly H -closed and strongly semiregular. We want to show that such a space X is regular because in a regular space we have that the closure and θ -closure are equal to each other and for this reason we have that the weakly H -closedness is equivalent to the H -closedness and we can apply Lemma 2.1. If X is strongly semiregular, then it has a base consisting of complements of closures of θ -open subsets of X which, by Proposition 2.3, are θ -open. So X has a base consisting of θ -open sets and for this reason it is regular. \square

We can observe that having a base consisting of complements of closures of θ -open sets is equivalent to having a base consisting of interiors of θ -closed sets.

We know that the character of a space is greater than or equal to the character of its semiregularization. We have the same result when considering the strongly semiregularization of the space X .

Proposition 2.5. *If X is a space and X_{ss} is the strong semiregularization of X , then $\chi(X_{ss}) \leq \chi(X)$.*

Proof. Let $x \in X$ and let \mathcal{B}_x be an open neighborhood system of x such that $|\mathcal{B}_x| \leq \chi(X)$. It suffices to show that $\{\text{int}([U]_\theta) : U \in \mathcal{B}_x\}$ is an open neighborhood system for x in X_{ss} . Let A be a θ -closed set such that $x \in \text{int}(A)$. There is $U \in \mathcal{B}_x$ such that $U \subseteq \text{int}(A) \subseteq A$. Thus, $x \in U \subseteq [U]_\theta \subseteq A$, and it follows that $x \in \text{int}([U]_\theta) \subseteq \text{int}(A)$. \square

3. CONSTRUCTION OF THE CARDINAL FUNCTION $\theta\text{-}aL'(\cdot)$ AND OF THE OPERATOR $\tilde{c}(\cdot)$

In this section we modify a filter construction given in [7]. In that paper an operator \hat{c} was constructed. Here we construct a related operator \tilde{c} .

Let X be a topological space, $x \in X$ and \mathcal{F}_x the collection of all finite intersections C of regular closed sets such that $x \in C$. It is easy to prove that \mathcal{F}_x is a filter base which can be extended to a filter \mathcal{C}_x that is maximal in the collection of all finite intersections of regular closed sets, partially ordered by inclusion.

The maximal filter \mathcal{C}_x has the following properties:

Proposition 3.1. *Let X be a topological space and $x \in X$. Every regular closed subset of X which meets every element of \mathcal{C}_x is an element of \mathcal{C}_x .*

Proof. Let U be an open subset of X and $x \in X$. \overline{U} is regular closed. Suppose \overline{U} meets every element of \mathcal{C}_x . Then $\{\overline{U}\} \cup \mathcal{C}_x$ is a filter base that contains \mathcal{C}_x which can be extended to a maximal filter \mathcal{M} . As \mathcal{C}_x is maximal, we have $\mathcal{C} = \mathcal{M}$ and $\overline{U} \in \mathcal{C}_x$. \square

Proposition 3.2. *Let X be a Urysohn space, then for every $x, y \in X$ with $x \neq y$ we have that $\mathcal{C}_x \neq \mathcal{C}_y$.*

Proof. Let $x, y \in X$ with $x \neq y$. Since X is a Urysohn space, there exist two open subsets U, V of X such that $x \in U, y \in V$ and $\overline{U} \cap \overline{V} = \emptyset$. We have that $\overline{U} \in \mathcal{C}_x$ and $\overline{V} \in \mathcal{C}_y$. If $\mathcal{C}_x = \mathcal{C}_y$, then $\overline{V} \in \mathcal{C}_x$ and $\overline{U} \cap \overline{V} = \emptyset \in \mathcal{C}_x$, a contradiction. \square

We now define new operators using the maximal filter \mathcal{C}_x .

Definition 3.1. For a space X and an open subset U of X , define:

$$\widetilde{U} = \{x \in X : \overline{U} \in \mathcal{C}_x\}.$$

In the following propositions we give several properties of \widetilde{U} .

Proposition 3.3. *Let X be a space and U an open subset of X . Then $\overline{U} \subseteq \widetilde{U} \subseteq \text{cl}_\theta(\overline{U})$.*

Proof. If $x \in \overline{U}$, then $\overline{U} \in \mathcal{C}_x$ and $x \in \widetilde{U}$. Let V be an open subset of X such that $x \in \overline{V}$. By Proposition 3.1, $\overline{V} \cap \overline{U} \neq \emptyset$, hence $x \in \text{cl}_\theta(\overline{U})$. \square

Question 3.1. If X is a space, does there exist an open subset U of X such that $\overline{U} \subsetneq \widetilde{U} \subsetneq \text{cl}_\theta(\overline{U})$?

Proposition 3.4. *If X is a topological space and V, W are open subsets of X , then $\widetilde{V} \cup \widetilde{W} = \widetilde{V \cup W}$. In particular this operator distributes over finite unions.*

Proof. $\widetilde{V} \cup \widetilde{W} = \{x \in X : \overline{V} \in \mathcal{C}_x\} \cup \{x \in X : \overline{W} \in \mathcal{C}_x\} = \{x \in X : \overline{V} \cup \overline{W} = \overline{V \cup W} \in \mathcal{C}_x\} = \widetilde{V \cup W}$. \square

The analogue of the following proposition in the case of Hausdorff spaces is contained in the proof of Proposition 4.1 in [7].

Proposition 3.5. *X is a Urysohn space if and only if for every $x, y \in X$ with $x \neq y$ there exist open subsets U, V of X such that $\widetilde{U} \cap \widetilde{V} = \emptyset$.*

Proof. Suppose that for every $x, y \in X$ with $x \neq y$ there exist open subsets U, V of X such that $\widetilde{U} \cap \widetilde{V} = \emptyset$. We have that $\overline{U} \subseteq \widetilde{U}$ and $\overline{V} \subseteq \widetilde{V}$, so $\overline{U} \cap \overline{V} = \emptyset$. This means that X is a Urysohn space.

Conversely, suppose X is Urysohn, then for every $x, y \in X$ with $x \neq y$ there exist open subsets U, V of X such that $\overline{U} \cap \overline{V} = \emptyset$. We want to show that $\widetilde{U} \cap \widetilde{V} = \emptyset$. In order to have a contradiction, suppose there exists $z \in \widetilde{U} \cap \widetilde{V}$. From the definition of \widetilde{U} and \widetilde{V} , we have $\overline{U}, \overline{V} \in \mathcal{C}_z$. \mathcal{C}_z is a filter, therefore $\overline{U} \cap \overline{V} = \emptyset \in \mathcal{C}_z$, a contradiction. \square

For a space X and a subset A of X we define a new operator called the \tilde{c} -closure in this way:

Definition 3.2. Let X be a space and A a subset of X ,

$$\tilde{c}(A) = \{x \in X : \widetilde{U} \cap A \neq \emptyset \text{ for every open subset } U \text{ of } X \text{ containing } x\}.$$

We say that $A \subseteq X$ is \tilde{c} -closed if $A = \tilde{c}(A)$.

We have the following propositions.

Proposition 3.6. *If X is a space and A is a subset of X , then we have $\text{cl}_\theta(A) \subseteq \tilde{c}(A) \subseteq \text{cl}_\gamma(A)$.*

Proof. If $x \in \text{cl}_\theta(A)$, then for every open subset V of X such that $x \in V$, $\overline{V} \cap A \neq \emptyset$. Therefore $\widetilde{V} \cap A \neq \emptyset$ and $x \in \tilde{c}(A)$.

If $x \in \tilde{c}(A)$, then for every open subset U of X such that $x \in U$ we have that $\widetilde{U} \cap A \neq \emptyset$ and by Proposition 3.3 we have $\widetilde{U} \subseteq \text{cl}_\theta(\overline{U})$. Thus for every open subset U of X such that $x \in U$ we have that $\text{cl}_\theta(\overline{U}) \cap A \neq \emptyset$, so $x \in \text{cl}_\gamma(A)$. \square

Question 3.2. If X is a space, does there exist $A \subseteq X$ such that $\text{cl}_\theta(A) \subsetneq \tilde{c}(A) \subsetneq \text{cl}_\gamma(A)$?

Proposition 3.7. *If X is a space and U is an open subset of X , then $\text{cl}_\theta(\overline{U}) \subseteq \tilde{c}(\overline{U}) \subseteq \tilde{c}(\widetilde{U})$.*

Proof. If U is an open subset of X , by definitions we have $\text{cl}_\theta(\overline{U}) \subseteq \tilde{c}(\overline{U}) \subseteq \tilde{c}(\widetilde{U})$. \square

Question 3.3. If X is a space, does there exist an open subset U of X such that $\text{cl}_\theta(\overline{U}) \subsetneq \tilde{c}(\widetilde{U})$?

We investigate the relation between Urysohn spaces and the operator $\tilde{c}(\cdot)$ and we prove the following:

Proposition 3.8. *If X is a Urysohn space, then for all $x, y \in X$ with $x \neq y$ there exists an open subset U of X such that $x \in U$ and $y \notin \tilde{c}(\widetilde{U})$.*

Proof. Let $x, y \in X$ with $x \neq y$. X is a Urysohn space, so that by Proposition 3.5 there exist open subsets U, V of X such that $x \in U$, $y \in V$ and $\widetilde{V} \cap \widetilde{U} = \emptyset$. We have that $\tilde{c}(\widetilde{U}) = \{z \in X : \widetilde{W} \cap \widetilde{U} \neq \emptyset \text{ for every open subset } U \text{ of } X \text{ and } y \in V \text{ but } \widetilde{V} \cap \widetilde{U} = \emptyset, \text{ so } y \notin \tilde{c}(\widetilde{U})\}$. \square

For a space X we define a new cardinal invariant $t_{\tilde{c}}(X)$ related to the tightness $t(X)$.

Definition 3.3. For a space X , the \tilde{c} -tightness of a point $x \in X$, denoted by $t_{\tilde{c}}(x, X)$, is

$$t_{\tilde{c}}(x, X) = \min\{k : \text{for every } A \subseteq X, \text{ if } x \in \tilde{c}(A), \text{ there exists } B \in [A]^{\leq k} \text{ such that } x \in \tilde{c}(B)\}.$$

The \tilde{c} -tightness of the space X , denoted by $t_{\tilde{c}}(X)$, is

$$t_{\tilde{c}}(X) = \sup_{x \in X} t_{\tilde{c}}(x, X).$$

A natural question is:

Question 3.4. Are $t_{\tilde{c}}(X)$ and $t(X)$ incomparable?

With the following proposition we prove that $t_{\tilde{c}}(X)$ is bounded above by the character.

Proposition 3.9. *If X is a space, then $t_{\tilde{c}}(X) \leq \chi(X)$.*

Proof. Let $x \in X$, $A \subseteq X$ such that $x \in \tilde{c}(A)$ and let \mathcal{V}_x be a neighborhood system of x in X with $|\mathcal{V}_x| \leq \chi(x, X)$. Because $x \in \tilde{c}(A)$, for every $V \in \mathcal{V}_x$ we have that $\tilde{V} \cap A \neq \emptyset$. Let $y_{\tilde{V}} \in \tilde{V} \cap A$ for every $V \in \mathcal{V}_x$. We put $B = \{y_{\tilde{V}} : V \in \mathcal{V}_x\}$, so $B \subseteq A$, $x \in \tilde{c}(B)$ and $|B| \leq \chi(x, X)$. This proves that $t_{\tilde{c}}(x, X) \leq \chi(x, X)$. \square

Using the \tilde{c} -tightness and the θ -pseudocharacter we find a bound for the cardinality of the \tilde{c} -closure of a subset A of a space X .

Proposition 3.10. *Let X be a Urysohn space such that $t_{\tilde{c}}(X)\psi_{\theta}(X) \leq k$. Then for every $A \subseteq X$ we have that $|\tilde{c}(A)| \leq |A|^k$.*

Proof. Let $x \in \tilde{c}(A)$. Since $\psi_{\theta}(X) \leq k$, by Proposition 3.8 there exists a family $\{U_{\alpha}(x)\}_{\alpha < k}$ of neighborhoods of x such that $\{x\} = \bigcap_{\alpha < k} \text{cl}_{\theta}(\overline{U_{\alpha}(x)}) = \bigcap_{\alpha < k} \tilde{c}(\overline{U_{\alpha}(x)})$. We want to prove that $x \in \tilde{c}(\overline{U_{\alpha}(x)} \cap A)$ for all $\alpha < k$. Let U be an open neighborhood of x and $\alpha < k$. Since $x \in \tilde{c}(A)$, we have that $\emptyset \neq \overline{U} \cap \overline{U_{\alpha}(x)} \cap A \subseteq \tilde{U} \cap \overline{U_{\alpha}(x)} \cap A$. This shows that $x \in \tilde{c}(\overline{U_{\alpha}(x)} \cap A)$. Since $t_{\tilde{c}}(X) \leq k$, there exists $A_{\alpha} \subset \overline{U_{\alpha}(x)} \cap A$ such that $|A_{\alpha}| \leq k$ and $x \in \tilde{c}(A_{\alpha}) \subseteq \tilde{c}(\overline{U_{\alpha}(x)})$. Then $\{x\} = \bigcap_{\alpha < k} \tilde{c}(A_{\alpha})$ and $\{A_{\alpha}\}_{\alpha < k} \in [[A]^{\leq k}]^{\leq k}$, so $|\tilde{c}(A)| \leq |[[A]^{\leq k}]^{\leq k}| = |A|^k$. \square

We now give another version of the Lindelöf and of the θ -almost Lindelöf degree using these new operators.

Definition 3.4. Let X be a topological space and Y a subset of X . We define $\tilde{L}(Y, X)$ by

$$\tilde{L}(Y, X) = \min \left\{ k : \text{for every cover } \mathcal{U} \text{ of } Y \text{ by sets open in } X \right. \\ \left. \text{there exists } \mathcal{V} \in [\mathcal{U}]^{\leq k} \text{ such that } X = \bigcup \tilde{\mathcal{V}} \right\}.$$

We put $\tilde{L}(X, X) = \tilde{L}(X)$.

We show now that if A is \tilde{c} -closed, then $\tilde{L}(A, X) \leq \tilde{L}(X)$.

Proposition 3.11. *If A is a \tilde{c} -closed subset of X , then $\tilde{L}(A, X) \leq \tilde{L}(X)$.*

Proof. Suppose that $\tilde{L}(X) \leq k$ and let C be a \tilde{c} -closed subset of X . For every $x \in X \setminus C$ there exists an open subset U_x of X such that $\tilde{U}_x \subseteq X \setminus C$. Let \mathcal{U} be an open cover of C , then $\mathcal{V} = \mathcal{U} \cup \{U_x : x \in X \setminus C\}$ is an open cover of X . As $\tilde{L}(X) \leq k$, there exists $\mathcal{V}' \in [\mathcal{V}]^{\leq k}$ such that $X = \bigcup \tilde{\mathcal{V}}'$. Thus there exists $\mathcal{V}'' \in [\mathcal{U}]^{\leq k}$ such that $C \subseteq \bigcup \tilde{\mathcal{V}}''$. This proves that $\tilde{L}(C, X) \leq k$. \square

Definition 3.5. For a space X , the θ -almost Lindelöf degree of X with respect to \hat{c} -closed subsets ($= \sup\{\theta\text{-}aL(C, X) : C \text{ is } \hat{c}\text{-closed}\}$) is denoted as $\theta\text{-}aL'(X)$ (instead of $\theta\text{-}aL_{\hat{c}}(X)$).

Note that $\theta\text{-}aL(X) \leq \theta\text{-}aL'(X) \leq \tilde{L}(X)$.

Question 3.5. Does there exist a space X such that $\theta\text{-}aL(X) < \theta\text{-}aL'(X) < \tilde{L}(X)$?

Now we prove our main result that is a new cardinal bound for Urysohn spaces. To prove this result we use Theorem 3.1 in [9].

Theorem 3.1 (Hodel). *Let X be a set, k an infinite cardinal, $f : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ an operator on X , and for each $x \in X$ let $\{V(\alpha, x) : \alpha < k\}$ be a collection of subsets of X . Assume*

- (T) (*tightness condition*) if $x \in f(H)$, then there exists $A \subseteq H$ with $|A| \leq k$ such that $x \in f(A)$;
- (C) (*cardinality condition*) if $A \subseteq X$ with $|A| \leq k$, then $|f(A)| \leq 2^k$;
- (C-S) (*cover-separation condition*) if $H \neq \emptyset$, $f(H) \subseteq H$ and $q \notin H$, then there exists $A \subseteq H$ with $|A| \leq k$ and a function $f : A \rightarrow k$ such that $H \subseteq \bigcup_{x \in A} V(f(x), x)$ and $q \notin \bigcup_{x \in A} V(f(x), x)$.

Then $|X| \leq 2^k$.

To prove the next theorem we use Theorem 3.1 and the operator $\tilde{c}(\cdot)$.

Theorem 3.2. *If X is a Urysohn space, then $|X| \leq 2^{\theta\text{-}aL'(X)t_{\tilde{c}}(X)\psi_{\theta}(X)}$.*

Proof. Let $k = \theta\text{-}aL'(X)t_{\tilde{c}}(X)\psi_{\theta}(X)$. As $\psi_{\theta}(X) \leq k$ for every $x \in X$ there exists a family $\mathcal{W}_x = \{W(\alpha, x) : \alpha < k\}$ of open subsets of X containing x such that $\{x\} = \bigcap_{W \in \mathcal{W}_x} \text{cl}_{\theta}(\overline{W})$.

For every $x \in X$ and $\alpha < k$, we put $V(\alpha, x) = \text{cl}_{\theta}(\overline{W(\alpha, x)})$ and prove the three conditions of Theorem 3.1.

For $H \subseteq X$, define $f(H) = \tilde{c}(H)$.

- ▷ Condition (T) is true because $t_{\tilde{c}} \leq k$.
- ▷ Condition (C) is true by Proposition 3.10.

▷ We prove condition (C-S). Let $\emptyset \neq H \subseteq X$ satisfy $\tilde{c}(H) \subseteq H$. We have that $H \subseteq \tilde{c}(H)$ so $H = \tilde{c}(H)$ and H is \tilde{c} -closed. Suppose $q \notin H$. For every $a \in H$ there exists $\alpha_a < k$ such that $q \notin \text{cl}_\theta(\overline{W(\alpha_a, a)}) = V(\alpha_a, a)$. Let $f: H \rightarrow X$ such that $f(a) = \alpha_a$. The set $\{W(f(a), a) : a \in H\}$ is an open cover of H and since H is \tilde{c} -closed and $\theta\text{-}aL'(X) \leq k$, there exists $A \in [H]^{\leq k}$ such that $H \subseteq \bigcup_{a \in A} V(f(a), a)$ and $q \notin \bigcup_{a \in A} V(f(a), a)$. This proves condition (C-S).

Applying Theorem 3.1 we have that $|X| \leq 2^k = 2^{\theta\text{-}aL'(X)t_{\tilde{c}}(X)\psi_\theta(X)}$. □

We can observe that every \tilde{c} -closed set is also θ -closed. We also know that the almost Lindelöf degree is hereditary with respect to θ -closed sets, so for every space X we have

$$\theta\text{-}aL'(X) \leq \theta\text{-}aL_\theta(X) \leq aL_\theta(X) = aL(X).$$

Furthermore, we know that $\psi_\theta(X) \leq \chi(X)$ and by Proposition 3.9 we have that $t_{\tilde{c}}(X) \leq \chi(X)$.

From these facts we obtain the following:

Corollary 3.1 ([4]). *If X is a Urysohn space, then $|X| \leq 2^{aL(X)\chi(X)}$.*

References

- [1] *O. T. Alas, L. D. Kočinac*: More cardinal inequalities on Urysohn spaces. *Math. Balk., New Ser.* 14 (2000), 247–251. zbl MR
- [2] *F. A. Basile, M. Bonanzinga, N. Carlson*: Variations on known and recent cardinality bounds. *Topology Appl.* 240 (2018), 228–237. zbl MR doi
- [3] *M. Bell, J. Ginsburg, G. Woods*: Cardinal inequalities for topological spaces involving the weak Lindelöf number. *Pac. J. Math.* 79 (1979), 37–45. zbl MR doi
- [4] *A. Bella, F. Cammaroto*: On the cardinality of Urysohn spaces. *Can. Math. Bull.* 31 (1988), 153–158. zbl MR doi
- [5] *F. Cammaroto, A. Catalioto, B. A. Pansera, B. Tsaban*: On the cardinality of the θ -closed hull of sets. *Topology Appl.* 160 (2013), 2371–2378. zbl MR doi
- [6] *F. Cammaroto, L. Kočinac*: On θ -tightness. *Facta Univ., Ser. Math. Inf.* 8 (1993), 77–85. zbl MR
- [7] *N. A. Carlson, J. R. Porter*: On the cardinality of Hausdorff spaces and H -closed spaces. *Topology Appl.* 241 (2018), 377–395. zbl MR doi
- [8] *R. Engelking*: *General Topology*. Wydawnictwo Naukowe PWN, Warsaw, 2007. (In Polish.) zbl
- [9] *R. E. Hodel*: Arhangel'skiĭ's solution to Alexandroff's problem: a survey. *Topology Appl.* 153 (2006), 2199–2217. zbl MR doi
- [10] *L. D. Kočinac*: On the cardinality of Urysohn and H -closed spaces. *Proc. Math. Conf., Priština, 1994* (L. D. Kočinac, ed.). Univ. of Priština, Faculty of Sciences, Priština, 1995, pp. 105–111. zbl MR
- [11] *L. D. Kočinac*: On the cardinality of Urysohn spaces. *Questions and Answers in General Topology* 13 (1995), 211–216. zbl MR
- [12] *J. R. Porter, R. G. Woods*: *Extensions and Absolutes of Hausdorff Spaces*. Springer, New York, 1988. zbl MR doi

- [13] *N. V. Veličko*: H -closed topological spaces. *Am. Math. Soc., Transl., II. Ser.* 78 (1968), 103–118; translated from *Mat. Sb., n. Ser.* 70 (1966), 98–112. [zbl](#) [MR](#) [doi](#)
- [14] *S. Willard, U. N. B. Dissanayake*: The almost Lindelöf degree. *Can. Math. Bull.* 27 (1984), 452–455. [zbl](#) [MR](#) [doi](#)

Authors' addresses: *Fortunata Aurora Basile*, University of Messina, Piazza Pugliatti, 1-98122 Messina, Italy, e-mail: basilef@unime.it; *Nathan Carlson*, Department of Mathematics, California Lutheran University, 60 W Olsen Rd, Thousand Oaks, CA 91360, California, USA, e-mail: ncarlson@callutheran.edu.