

## AN OBSERVATION ON SPACES WITH A ZEROSET DIAGONAL

WEI-FENG XUAN, Nanjing

Received February 6, 2018. Published online November 26, 2018.

Communicated by Pavel Pyrih

*Abstract.* We say that a space  $X$  has the discrete countable chain condition (DCCC for short) if every discrete family of nonempty open subsets of  $X$  is countable. A space  $X$  has a zerset diagonal if there is a continuous mapping  $f: X^2 \rightarrow [0, 1]$  with  $\Delta_X = f^{-1}(0)$ , where  $\Delta_X = \{(x, x): x \in X\}$ . In this paper, we prove that every first countable DCCC space with a zerset diagonal has cardinality at most  $\mathfrak{c}$ .

*Keywords:* first countable; discrete countable chain condition; zerset diagonal; cardinal

*MSC 2010:* 54D20, 54E35

## 1. INTRODUCTION

All topological spaces in this paper are assumed to be Hausdorff unless otherwise stated. The cardinality of a set  $X$  is denoted by  $|X|$ , and  $[X]^2$  will denote the set of two-element subsets of  $X$ . We write  $\omega$  for the first infinite cardinal,  $\omega_1$  for the first uncountable cardinal and  $\mathfrak{c}$  for the cardinality of the continuum.

In 1977, Ginsburg and Woods proved that the cardinality of a  $T_1$ -space with countable extent and a  $G_\delta$ -diagonal is at most  $\mathfrak{c}$  (see [5]). In the same paper, Ginsburg and Woods asked if it was true that a regular CCC-space (here CCC denotes the countable chain condition) with a  $G_\delta$ -diagonal has cardinality at most  $\mathfrak{c}$ . This question was also posted by Arhangel'skii independently. In 1984, Shakhmatov showed that cardinalities of such spaces may not have an upper bound (see [8]). And later, Uspenskij proved that an upper bound still does not exist even assuming Fréchet property (see [9]). Regular  $G_\delta$ -diagonal is a property stronger than  $G_\delta$ -diagonal. Arhangel'skii asked what if “ $G_\delta$ -diagonal” is replaced by “regular  $G_\delta$ -diagonal”.

---

The author is supposed by NSFC Projects 11801271 and 11626131.

In 2005, Buzyakova proved that the cardinality of a CCC-space with a regular  $G_\delta$ -diagonal is at most  $\mathfrak{c}$  (see [3]). In 2015, Gotchev in [6] proved that the cardinality of a weakly Lindelöf space with a regular  $G_\delta$ -diagonal is at most  $2^\mathfrak{c}$ .

**Definition 1.1.** We say that a space  $X$  has the *discrete countable chain condition* (DCCC for short) if every discrete family of nonempty open subsets of  $X$  is countable.

By Definition 1.1, it follows immediately that every CCC space is DCCC. In fact, every weakly Lindelöf space is DCCC, but the converse is not true. For example,  $\omega_1$  with the ordered topology is a first countable and countably compact (hence, DCCC) space which is not weakly Lindelöf, because the open cover  $\mathcal{U} = \{[0, \alpha] : \alpha < \omega_1\}$  of  $\omega_1$  does not have a countable subfamily whose union is dense in  $\omega_1$ .

**Definition 1.2** ([2]). A space  $X$  has a *zeroset diagonal* if there is a continuous mapping  $f: X^2 \rightarrow [0, 1]$  with  $\Delta_X = f^{-1}(0)$ , where  $\Delta_X = \{(x, x) : x \in X\}$ .

It is well-known and easy to prove that every submetrizable space has a zeroset diagonal and every zeroset diagonal is a regular  $G_\delta$ -diagonal. The converses are not true (see [1], [10]).

In this paper, we prove that every first countable DCCC space with a zeroset diagonal has cardinality at most  $\mathfrak{c}$ .

All notations and terminology not explained in the paper are given in [4].

## 2. RESULTS

We will use the following countable version of a set-theoretic theorem due to Erdős and Radó (see [7], page 8).

**Lemma 2.1.** *Let  $X$  be a set with  $|X| > \mathfrak{c}$  and suppose  $[X]^2 = \bigcup \{P_n : n \in \omega\}$ . Then there exist  $n_0 < \omega$  and a subset  $S$  of  $X$  with  $|S| > \omega$  such that  $[S]^2 \subset P_{n_0}$ .*

**Theorem 2.2.** *Every first countable DCCC space  $X$  with a zeroset diagonal has cardinality at most  $\mathfrak{c}$ .*

**Proof.** Assume the contrary, i.e. that  $|X| > \mathfrak{c}$ . Fix a continuous function  $f: X^2 \rightarrow [0, 1]$  with  $\Delta_X = f^{-1}(0)$ . Let  $\mathcal{B}(x) = \{B_n(x) : n \in \omega\}$  be a local decreasing base for each  $x \in X$ . Since for any distinct  $x, y \in X$  there is some  $n_1 \in \omega$  such that  $(x, y) \in f^{-1}((1/(n_1 + 2019), 1])$  and since  $f$  is continuous, there are  $n_2, n_3 \in \omega$  such that

$$B_{n_2}(x) \times B_{n_3}(y) \subset f^{-1}\left(\left(\frac{1}{n_1 + 2019}, 1\right]\right).$$

Let  $n^* = \max\{n_1, n_2, n_3\}$ . Then by our hypothesis, we can deduce that

$$B_{n^*}(x) \times B_{n^*}(y) \subset f^{-1}\left(\left(\frac{1}{n^* + 2019}, 1\right]\right).$$

Thus, the following sets  $P_n$  are well defined. For each  $n \in \omega$  let

$$P_n = \left\{ \{x, y\} \in [X]^2 : B_n(x) \times B_n(y) \subset f^{-1}\left(\left(\frac{1}{n + 2019}, 1\right]\right) \right\}.$$

It is clear that  $[X]^2 = \bigcup \{P_n : n \in \omega\}$ . (Note that  $[X]^2$  is the set of two-element subsets of  $X$ ). We can apply Lemma 2.1 to conclude that there exists an uncountable subset  $S$  of  $X$  and  $n_0 \in \omega$  such that  $[S]^2 \subset P_{n_0}$ . It follows immediately that  $\mathcal{U} = \{B_{n_0}(x) : x \in S\}$  is an uncountable family of nonempty open sets of  $X$ . Since  $X$  is DCCC, the family  $\mathcal{U}$  must have a cluster point  $x \in X$ . Pick any neighbourhood  $O_x$  of  $x$  such that

$$O_x \times O_x \subset f^{-1}\left(\left[0, \frac{1}{n_0 + 2019}\right)\right).$$

Obviously,  $O_x$  meets infinitely many members of  $\mathcal{U}$ . Thus, there exist two distinct (at least)  $y, z \in S$  such that  $O_x \cap B_{n_0}(y) \neq \emptyset$  and  $O_x \cap B_{n_0}(z) \neq \emptyset$ . Take any  $y' \in O_x \cap B_{n_0}(y)$  and  $z' \in O_x \cap B_{n_0}(z)$ . Hence,  $f(y', z') < 1/(n_0 + 2019)$  since  $y', z' \in O_x$ . On the other hand,  $f(y', z') > 1/(n_0 + 2019)$  since  $y' \in B_{n_0}(y)$ ,  $z' \in B_{n_0}(z)$  and  $\{y, z\} \in P_{n_0}$ . This gives a contradiction and we prove that  $|X| \leq \mathfrak{c}$ .  $\square$

If we drop the condition “DCCC”, or “zeroset diagonal” in Theorem 2.2, the conclusion is no longer true, which can be seen in the following examples.

**Example 2.3.** Let  $D$  be a discrete space with  $|D| = 2^{\mathfrak{c}}$ . It is evident that  $D$  is first countable and has a zeroset diagonal, but  $D$  is not DCCC.

**Example 2.4.** Let  $X$  be the subspace of  $[0, 2^{\mathfrak{c}}]$ , consisting of all ordinals of countable cofinality, equipped with the ordered topology. Then  $X$  is a first countable and countably compact (hence DCCC) space of cardinality  $2^{\mathfrak{c}}$ , but it does not have a zeroset diagonal.

We finish the paper with the following question.

**Question 2.5.** Is it true that every DCCC (or weakly Lindelöf) space with a zeroset diagonal has cardinality at most  $\mathfrak{c}$ ?

**Acknowledgement.** We would like to thank the referee for their valuable remarks and suggestions which greatly improved the paper.

## References

- [1] *A. V. Arhangel'skii, R. Z. Buzyakova*: The rank of the diagonal and submetrizability. *Commentat. Math. Univ. Carol.* **47** (2006), 585–597. zbl MR
- [2] *R. Z. Buzyakova*: Observations on spaces with zeroset or regular  $G_\delta$ -diagonals. *Commentat. Math. Univ. Carol.* **46** (2005), 469–473. zbl MR
- [3] *R. Z. Buzyakova*: Cardinalities of ccc-spaces with regular  $G_\delta$ -diagonals. *Topology Appl.* **153** (2006), 1696–1698. zbl MR doi
- [4] *R. Engelking*: *General Topology*. Sigma Series in Pure Mathematics 6. Heldermann, Berlin, 1989. zbl MR
- [5] *J. Ginsburg, R. G. Woods*: A cardinal inequality for topological spaces involving closed discrete sets. *Proc. Am. Math. Soc.* **64** (1977), 357–360. zbl MR doi
- [6] *I. S. Gotchev*: Cardinalities of weakly Lindelöf spaces with regular  $G_\kappa$ -diagonals. Available at <https://arxiv.org/abs/1504.01785> (2015).
- [7] *R. E. Hodel*: Cardinal function. I. *Handbook of Set-Theoretic Topology* (K. Kunen et al., eds.). North-Holland, Amsterdam, 1984, pp. 1–61. zbl MR doi
- [8] *D. Shakhmatov*: No upper bound for cardinalities of Tychonoff c.c.c. spaces with a  $G_\delta$ -diagonal exists. An answer to J. Ginsburg and R. G. Woods' question. *Commentat. Math. Univ. Carol.* **25** (1984), 731–746. zbl MR
- [9] *V. V. Uspenskij*: A large  $F_\sigma$ -discrete Fréchet space having the Souslin property. *Commentat. Math. Univ. Carol.* **25** (1984), 257–260. zbl MR
- [10] *M. L. Wage, W. G. Fleissner, G. M. Reed*: Normality versus countable paracompactness in perfect spaces. *Bull. Am. Math. Soc.* **82** (1976), 635–639. zbl MR doi

*Author's address:* Wei-Feng Xuan, School of Statistics and Mathematics, Nanjing Audit University, 86 West Yushan Road, Pukou District, Nanjing, 211815, China, e-mail: [wfxuan@nau.edu.cn](mailto:wfxuan@nau.edu.cn).