

SOME RELATIONS ON HUMBERT MATRIX POLYNOMIALS

AYMAN SHEHATA, Assiut, Qassim

Received February 18, 2014. First published August 2, 2016.

Communicated by Dagmar Medková

Abstract. The Humbert matrix polynomials were first studied by Khammash and Shehata (2012). Our goal is to derive some of their basic relations involving the Humbert matrix polynomials and then study several generating matrix functions, hypergeometric matrix representations, matrix differential equation and expansions in series of some relatively more familiar matrix polynomials of Legendre, Gegenbauer, Hermite, Laguerre and modified Laguerre. Finally, some definitions of generalized Humbert matrix polynomials also of two, three and several index are derived.

Keywords: hypergeometric matrix function; Humbert matrix polynomials; matrix functional calculus; generating matrix function; matrix differential equation

MSC 2010: 15A60, 33C55, 33C45, 33E20

1. INTRODUCTION AND PRELIMINARIES

Special matrix functions are very close to statistics, Lie group theory and number theory [14]. In the recent work, matrix polynomials have significant emergent in [1]–[7], [9], [10], [12], [17]–[21], [24]–[32], [34]–[37]. Results in the theory of classical orthogonal polynomials have been extended to orthogonal matrix polynomials in [8], [16], [38]. Humbert matrix polynomials have been introduced and studied in [22]. The reasons of interest in this family of Humbert matrix polynomials are their intrinsic mathematical importance.

Our main aim in this paper is to prove new properties the generalized hypergeometric matrix function and Humbert matrix polynomials. The structure of this paper is as follows. In Section 2, some basic relations involving the generalized hypergeometric matrix functions and convergence properties, integral representation, differential properties, and matrix differential equation are obtained. In Section 3, we give some new results involving the Humbert matrix polynomials and then establish

several operational results, matrix differential equation, finite series representation, hypergeometric representations, four additional generating matrix functions and expansions of Humbert matrix polynomials in series of other polynomials which are best stated in terms of the generalized matrix polynomials. Relationships with other polynomial systems are also developed. Finally, we define the multiindex Humbert matrix polynomials and present several new results for matrix polynomials in Section 4.

Throughout this paper, for a matrix A in $\mathbb{C}^{N \times N}$, its spectrum $\sigma(A)$ denotes the set of all eigenvalues of A . Its two-norm will be denoted by $\|A\|$, and is defined by

$$\|A\| = \sup_{x \neq 0} \frac{\|Ax\|_2}{\|x\|_2}$$

where, for a vector $x \in \mathbb{C}^N$, $\|x\|_2 = (x^T x)^{1/2}$ is the Euclidean norm of x .

If $f(z)$ and $g(z)$ are holomorphic functions of the complex variable z , which are defined in an open set Ω of the complex plane, and A, B are matrices in $\mathbb{C}^{N \times N}$ with $\sigma(A) \subset \Omega$ and $\sigma(B) \subset \Omega$, such that $AB = BA$, then from the properties of the matrix functional calculus in [13] it follows that

$$(1.1) \quad f(A)g(B) = g(B)f(A).$$

We recall that the reciprocal gamma function denoted by $\Gamma^{-1}(z) = 1/\Gamma(z)$ is an entire function of the complex variable z and thus for any matrix A in $\mathbb{C}^{N \times N}$, $\Gamma^{-1}(A)$ is a well defined matrix. Furthermore, if A is a matrix in $\mathbb{C}^{N \times N}$ such that

$$(1.2) \quad A + nI \quad \text{is an invertible matrix for all integers } n \geq 0$$

where I is the identity matrix in $\mathbb{C}^{N \times N}$, then the Pochhammer symbol or shifted factorial is defined by [15]

$$(1.3) \quad \begin{aligned} (A)_n &= A(A+I)(A+2I) \dots (A+(n-1)I) \\ &= \Gamma(A+nI)\Gamma^{-1}(A); \quad n \geq 1, (A)_0 = I. \end{aligned}$$

For matrices $A(k, n)$ and $B(k, n)$ in $\mathbb{C}^{N \times N}$ where $n \geq 0, k \geq 0$ the following relations are satisfied according to Defez and Jódar in [11]:

$$(1.4) \quad \begin{aligned} \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} B(k, n) &= \sum_{n=0}^{\infty} \sum_{k=0}^n B(k, n-k), \\ \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} A(k, n) &= \sum_{n=0}^{\infty} \sum_{k=0}^{[n/m]} A(k, n-mk); \quad m \in \mathbb{N}. \end{aligned}$$

Similarly, we can write

$$(1.5) \quad \sum_{n=0}^{\infty} \sum_{k=0}^n B(k, n) = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} B(k, n+k),$$

$$\sum_{n=0}^{\infty} \sum_{k=0}^{[n/m]} A(k, n) = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} A(k, n+mk); \quad m \in \mathbb{N}.$$

Let us denote real numbers $M(A)$ and $m(A)$ by

$$(1.6) \quad M(A) = \max\{\Re(z) : z \in \sigma(A)\}; \quad m(A) = \min\{\Re(z) : z \in \sigma(A)\}.$$

Note that, if k is large enough so that for $k > \|C\|$, then we have the relation which is given in Jódar and Cortés [18], [17] in the form

$$(1.7) \quad \|(C + kI)^{-1}\| \leq \frac{1}{k - \|C\|}; \quad k > \|C\|.$$

The hypergeometric matrix function ${}_2F_1(A, B; C; z)$ has been given in [17] in the form

$$(1.8) \quad {}_2F_1(A, B; C; z) = \sum_{k=0}^{\infty} \frac{(A)_k (B)_k ((C)_k)^{-1}}{k!} z^k$$

where A, B and C are matrices of $\mathbb{C}^{N \times N}$ such that $C + nI$ is an invertible matrix for all integers $n \geq 0$. It has been shown by Jódar and Cortés [17] that the series is absolutely convergent for $|z| = 1$ when

$$m(C) > M(A) + M(B).$$

If $CB = BC$ and $C, B, C - B$ are positive stable matrices, then, for $|z| < 1$, an integral representation of (1.8) was given in the form [17]

$$(1.9) \quad {}_2F_1(A, B; C; z) = \int_0^1 (1-tz)^{-A} t^{B-I} (1-t)^{C-B-I} dt$$

$$\times \Gamma^{-1}(B) \Gamma^{-1}(C-B) \Gamma(C).$$

For any matrix A in $\mathbb{C}^{N \times N}$, we will exploit the relation due to [17]

$$(1.10) \quad (1-z)^{-A} = {}_1F_0(A; -; z) = \sum_{n=0}^{\infty} \frac{1}{n!} (A)_n z^n; \quad |z| < 1.$$

Our main aim in the next section is to prove new properties of generalized hypergeometric matrix function.

2. PROPERTIES OF GENERALIZED HYPERGEOMETRIC MATRIX FUNCTION

The generalized hypergeometric matrix function is defined as follows:

Definition 2.1. Let A_1, A_2, A_3, B_1 and B_2 be matrices in $\mathbb{C}^{N \times N}$ such that

$$(2.1) \quad B_1 + nI \text{ and } B_2 + nI \text{ are invertible matrices for all integers } n \geq 0.$$

Then we define the generalized hypergeometric matrix function as

$$(2.2) \quad {}_3F_2(A_1, A_2, A_3; B_1, B_2; z) = \sum_{k=0}^{\infty} \frac{z^k}{k!} (A_1)_k (A_2)_k (A_3)_k ((B_1)_k)^{-1} ((B_2)_k)^{-1}.$$

We are going to study the convergence properties of the hypergeometric matrix function. Note that if k is large enough so that $k > \|B_j\|$; $j = 1, 2$, then by the perturbation lemma, see [17], we can write

$$\left\| \left(\frac{1}{k} B_j + I \right)^{-1} \right\| \leq \frac{1}{1 - k^{-1} \|B_j\|} = \frac{k}{k - \|B_j\|}; \quad j = 1, 2$$

and

$$(2.3) \quad \begin{aligned} \|(B_j + kI)^{-1}\| &= \left\| \frac{1}{k} \left(\frac{1}{k} B_j + I \right)^{-1} \right\| = \frac{1}{k} \left\| \left(\frac{1}{k} B_j + I \right)^{-1} \right\| \\ &\leq \frac{1}{k - \|B_j\|}; \quad k > \|B_j\|; \quad j = 1, 2. \end{aligned}$$

Let us denote

$$(2.4) \quad T_j(k) = \|B_j^{-1}\| \|(B_j + I)^{-1}\| \dots \|(B_j + (k-1)I)^{-1}\|$$

for $k \geq 1$ and $j = 1, 2$.

Note that from (1.3) we obtain

$$(2.5) \quad \|(A_i)_k\| \leq (\|A_i\|)_k; \quad i = 1, 2, 3.$$

Using (2.3), (2.4) and (2.5) for $k > \|B_j\|$ we have

$$(2.6) \quad \begin{aligned} \left\| \frac{z^k}{k!} (A_1)_k (A_2)_k (A_3)_k ((B_1)_k)^{-1} ((B_2)_k)^{-1} \right\| \\ \leq \frac{1}{k!} |z^k| \| (A_1)_k \| \| (A_2)_k \| \| (A_3)_k \| T_1(k) T_2(k) \\ \leq \frac{1}{k!} |z^k| (\|A_1\|)_k (\|A_2\|)_k (\|A_3\|)_k T_1(k) T_2(k). \end{aligned}$$

Now we will investigate the convergence of the series

$$(2.7) \quad \sum_{k=0}^{\infty} \frac{1}{k!} |z^k| (\|A_1\|)_k (\|A_2\|)_k (\|A_3\|)_k T_1(k) T_2(k).$$

Using the ratio test and the relation (1.7), one gets

$$(2.8) \quad \lim_{k \rightarrow \infty} \left| \frac{(\|A_1\|)_{k+1} (\|A_2\|)_{k+1} (\|A_3\|)_{k+1} T_1(k+1) T_2(k+1) k! z^{k+1}}{(\|A_1\|)_k (\|A_2\|)_k (\|A_3\|)_k T_1(k) T_2(k) (k+1)! z^k} \right| \\ \leq \lim_{k \rightarrow \infty} \left| \frac{(\|A_1\| + k) (\|A_2\| + k) (\|A_3\| + k) \| (B_1 + kI)^{-1} \| \| (B_2 + kI)^{-1} \| z^{k+1}}{(k+1) z^k} \right| \\ \leq \lim_{k \rightarrow \infty} \frac{(\|A_1\| + k) (\|A_2\| + k) (\|A_3\| + k)}{(k - \|B_1\|) (k - \|B_2\|) (k+1)} |z|,$$

hence the power series (2.2) is absolutely convergent for $|z| < 1$ and diverges for $|z| > 1$. Analogously to Theorem 3 in [17], we can state

Theorem 2.1. *The generalized hypergeometric matrix function ${}_3F_2(A_1, A_2, A_3; B_1, B_2; z)$ is absolutely convergent for $|z| = 1$ when*

$$(2.9) \quad m(B_1) + m(B_2) > M(A_1) + M(A_2) + M(A_3)$$

where M and m are defined in (1.6).

The integral representation (1.9) of the generalized hypergeometric matrix function can be extended to obtain the following result:

Theorem 2.2. *Let A_i , $1 \leq i \leq 3$ and B_j , $1 \leq j \leq 2$ be matrices in $\mathbb{C}^{N \times N}$ such that $B_j + kI$ are invertible matrices for all integers $k \geq 0$. Suppose that the matrices A_i , B_j and $B_j - A_i$ are positive stable matrices. For $|z| < 1$ and $|zt| < 1$ it follows that*

$$(2.10) \quad {}_3F_2(A_1, A_2, A_3; B_1, B_2; z) = \int_0^1 t^{A_1 - I} (1-t)^{B_1 - A_1 - I} {}_2F_1(A_2, A_3; B_2; zt) dt \\ \times \Gamma^{-1}(A_1) \Gamma^{-1}(B_1 - A_1) \Gamma(B_1)$$

where $A_i B_j = B_j A_i$.

The proof is similar to the proof of Theorem 5 in [17].

Consider the differential operator $\theta = z(d/dz)$, $D_z = d/dz$, $\theta z^k = kz^k$. For commutative matrices, we have

$$\begin{aligned} & \theta(\theta I + B_1 - I)(\theta I + B_2 - I)_3F_2 \\ &= \sum_{k=1}^{\infty} \frac{kz^k}{k!} (kI + B_1 - I)(kI + B_2 - I)(A_1)_k(A_2)_k(A_3)_k((B_1)_k)^{-1}((B_2)_k)^{-1} \\ &= \sum_{k=1}^{\infty} \frac{z^k}{(k-1)!} (A_1)_k(A_2)_k(A_3)_k((B_1)_{k-1})^{-1}((B_2)_{k-1})^{-1}. \end{aligned}$$

Replacing k by $k+1$, we have

$$\begin{aligned} & \theta(\theta I + B_1 - I)(\theta I + B_2 - I)_3F_2 \\ &= \sum_{k=0}^{\infty} \frac{z^{k+1}}{k!} (A_1)_{k+1}(A_2)_{k+1}(A_3)_{k+1}((B_1)_k)^{-1}((B_2)_k)^{-1} \\ &= z(\theta I + A_1)(\theta I + A_2)(\theta I + A_3)_3F_2. \end{aligned}$$

Thus, we have shown that ${}_3F_2$ is a solution of the matrix differential equation

$$(\theta(\theta I + B_1 - I)(\theta I + B_2 - I) - z(\theta I + A_1)(\theta I + A_2)(\theta I + A_3))_3F_2 = \mathbf{0}$$

where $\mathbf{0}$ is a zero matrix or null matrix in $\mathbb{C}^{N \times N}$. This result is summarized below.

Theorem 2.3. *Let A_1, A_2, A_3, B_1 and B_2 be matrices in $\mathbb{C}^{N \times N}$ satisfying the spectral condition (2.1) and $B_j A_i = A_i B_j$; $1 \leq i \leq 3$ and $1 \leq j \leq 2$. Then ${}_3F_2$ satisfies the matrix differential equation*

$$(2.11) \quad (\theta(\theta I + B_1 - I)(\theta I + B_2 - I) - z(\theta I + A_1)(\theta I + A_2)(\theta I + A_3))_3F_2 = \mathbf{0}$$

or

$$\begin{aligned} (2.12) \quad & \left((1-z)z^2 \frac{d^3}{dz^3} + ((B_1 + B_2 + I) - (A_1 + A_2 + A_3 + 3)z)z \frac{d^2}{dz^2} \right. \\ & \left. + (B_1 B_2 - z(A_1 A_2 + A_2 A_3 + A_1 A_3 + A_1 + A_2 + A_3 + I)) \frac{d}{dz} \right. \\ & \left. - A_1 A_2 A_3 \right)_3F_2 = \mathbf{0}. \end{aligned}$$

3. PROPERTIES OF HUMBERT MATRIX POLYNOMIALS

In this section, we deal with the Humbert matrix polynomials $h_n^A(x)$ that are defined by (see [22])

$$(3.1) \quad h_n^A(x) = \sum_{k=0}^{[n/3]} \frac{(-1)^k (3x)^{n-3k}}{k!(n-3k)!} (A)_{n-2k}$$

for a matrix A in $\mathbb{C}^{N \times N}$ such that

$$(3.2) \quad A + nI \text{ is an invertible matrix for all integers } n \geq 0.$$

Such matrix polynomials have the generating matrix function

$$(3.3) \quad (1 - 3xt + t^3)^{-A} = \sum_{n=0}^{\infty} h_n^A(x) t^n; \quad |3xt - t^3| < 1,$$

and the hypergeometric matrix representation for Humbert matrix polynomials

$$(3.4) \quad h_n^A(x) = \frac{(A)_n (3x)^n}{n!} {}_3F_2 \left(-\frac{1}{3}nI, \frac{1}{3}(1-n)I, \frac{1}{3}(2-n)I; \right. \\ \left. \frac{1}{2}(I - A - nI), \frac{1}{2}(2I - A - nI); \frac{1}{4x^3} \right)$$

where $(I - A - nI)/2$ and $(2I - A - nI)/2$ are matrices in $\mathbb{C}^{N \times N}$ satisfying the condition (2.1).

Next, let us recall the convergent properties and matrix differential equations of Humbert matrix polynomials which will be used in the following.

Theorem 3.1. *Let A be a matrix in $\mathbb{C}^{N \times N}$ satisfying the spectral condition (3.2). Then the power series (3.1) is absolutely convergent for $|1/4x^3| < 1$ and diverges for $|1/4x^3| > 1$.*

Theorem 3.2. *Let A be a matrix in $\mathbb{C}^{N \times N}$ satisfying the condition (3.2). Then Humbert matrix polynomials are absolutely convergent for $|1/4x^3| = 1$ when*

$$(3.5) \quad m \left(\frac{1}{2}(2I - A - nI) \right) + m \left(\frac{1}{2}(I - A - nI) \right) \\ > M \left(-\frac{1}{3}nI \right) + M \left(\frac{1}{3}(1-n)I \right) + M \left(\frac{1}{3}(2-n)I \right)$$

where M and m are defined in (1.6).

From the fundamental relation (3.3) by the usual method in such cases (differentiating (3.3) with respect to x and then with respect to t) the following recurrence formula may be easily obtained:

$$(3.6) \quad (n+1)h_{n+1}^A(x) - 3x(A+nI)h_n^A(x) + (3A+(n-2)I)h_{n-2}^A(x) = \mathbf{0}, \quad n \geq 2,$$

see [22]. This formula affects only Humbert matrix polynomials with the same superior index. But another one can be readily written, connecting Humbert matrix polynomials with different indices, both superior and inferior, namely

$$(3.7) \quad (n+1)h_{n+1}^A(x) + 3Ah_{n-2}^{A+I}(x) = 3Axh_n^{A+I}(x), \quad n \geq 2.$$

In [22], other recurrence formulae introduce the differential coefficient of the Humbert matrix polynomials:

$$(3.8) \quad nh_n^A(x) + \frac{d}{dx}h_{n-2}^A(x) - x\frac{d}{dx}h_n^A(x) = \mathbf{0}, \quad n \geq 2,$$

$$(3.9) \quad (3A+nI)h_n^A(x) = \frac{d}{dx}h_{n+1}^A(x) - 2x\frac{d}{dx}h_n^A(x)$$

and

$$(3.10) \quad \frac{d}{dx}h_n^A(x) = 3Ah_{n-1}^{A+I}(x), \quad n \geq 1.$$

With help of these recurrence formulas, we can write the following equations: The first of them is

$$(3.11) \quad \frac{d}{dx}h_{n+1}^A(x) = 3Ah_n^{A+I}(x).$$

Comparing (3.11) with (3.10), we obtain

$$(3.12) \quad \frac{d^2}{dx^2}h_{n+2}^A(x) = 9A(A+I)h_n^{A+2I}(x).$$

Due to (3.12), it is easy to see that

$$(3.13) \quad D^k h_{n+k}^A(x) = 3^k(A)_k h_n^{A+kI}(x); \quad D = \frac{d}{dx}.$$

From the relations (3.8) and (3.9), we easily get the relations

$$(3.14) \quad (xD-n)h_n^A(x) = Dh_{n-2}^A(x), \quad n \geq 2$$

and

$$(3.15) \quad (2xDI + nI + 3A)h_n^A(x) = Dh_{n+1}^A(x).$$

Now iteratively applying the linear differential operator in (3.15) twice to (3.14) and using the simple relation

$$(3.16) \quad (axD + b)D^n = D^n(axD + b - na)$$

the matrix differential equation is given in the form

$$(3.17) \quad (2xDI + (n + 3)I + 3A)(2xDI + nI + 3A)(xD - n)h_n^A(x) = D^3h_n^A(x).$$

From (3.17), we get the matrix differential equation of third order satisfied by Humbert matrix polynomials

$$(3.18) \quad \begin{aligned} & (1 - 4x^3)D^3h_n^A(x) - 6(3I + 2A)x^2D^2h_n^A(x) \\ & - (((n + 5)I + 3A)((2 - 3n)I + 3A) + 10nI)xDh_n^A(x) \\ & + n(nI + 3A)((n + 3)I + 3A)h_n^A(x) = \mathbf{0}. \end{aligned}$$

This result is summarized in

Theorem 3.3. *Let A be a matrix in $\mathbb{C}^{N \times N}$ satisfying the spectral condition (3.2). Then the Humbert matrix polynomial is the solution of the matrix differential equation (3.18).*

A variety of interesting results can be easily deduced from Theorem 2.3. In the next result, the Humbert matrix polynomials appear as finite series solutions of the third order matrix differential equation.

Corollary 3.1. *The Humbert matrix polynomial is the solution of the matrix differential equation of the third order*

$$(3.19) \quad \begin{aligned} & (1 - 4x^3)D^3h_n^A(x) - 6(3I + 2A)x^2D^2h_n^A(x) \\ & - (((n + 5)I + 3A)((2 - 3n)I + 3A) + 10nI)xDh_n^A(x) \\ & + n(nI + 3A)((n + 3)I + 3A)h_n^A(x) = \mathbf{0}. \end{aligned}$$

Proof. Put $z = 1/4x^3$, $A_1 = -nI/3$, $A_2 = (1 - n)I/3$, $A_3 = (2 - n)I/3$, $B_1 = (I - A - nI)/2$ and $B_2 = (2I - A - nI)/2$ in (2.12) and multiply by $(3x)^n$, then the proof of (3.19) follows directly. \square

Now we obtain the finite series representation for Humbert matrix polynomials $h_n^A(x)$.

Theorem 3.4. Let A be a matrix in $\mathbb{C}^{N \times N}$ satisfying (3.2). Then we have

$$(3.20) \quad h_n^A(x) = \sum_{k=0}^{[n/2]} \sum_{s=0}^{n-2k} \frac{(A)_k(-k)_s(2A+2kI)_{n-2k-s} \left(\frac{3}{2}x\right)^{n-3s}}{k!s!(n-2k-s)!}.$$

Proof. From (3.3) and (1.5) we have

$$(3.21) \quad \begin{aligned} \sum_{n=0}^{\infty} h_n^A(x)t^n &= (1-3xt+t^3)^{-A} = \left(\left(1 - \frac{3xt}{2}\right)^2 - \left(\frac{3xt}{2}\right)^2 + t^3 \right)^{-A} \\ &= \left(1 - \frac{3xt}{2}\right)^{-2A} \left(1 - \frac{(3xt)^2 2^{-2} - t^3}{(1-3xt2^{-1})^2}\right)^{-A} \\ &= \left(1 - \frac{3xt}{2}\right)^{-2A} \sum_{k=0}^{\infty} \frac{(A)_k}{k!} \left(\frac{(3xt)^2 2^{-2} - t^3}{(1-3xt2^{-1})^2}\right)^k \\ &= \sum_{k=0}^{\infty} \frac{(A)_k}{k!} \left(\left(\frac{3xt}{2}\right)^2 - t^3\right)^k \left(1 - \frac{3xt}{2}\right)^{-2A-2kI} \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{(A)_k(2A+2kI)_n \left(\frac{3}{2}xt\right)^{n+2k}}{k!n!} \left(1 - \frac{4t}{(3x)^2}\right)^k \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \sum_{s=0}^k \frac{(A)_k(-k)_s(2A+2kI)_n \left(\frac{3}{2}x\right)^{n+2k-2s}}{k!s!n!} t^{n+2k+s} \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \sum_{s=0}^{\infty} \frac{(A)_k(-k)_s(2A+2kI)_n \left(\frac{3}{2}x\right)^{n+2k-2s}}{k!s!n!} t^{n+2k+s}. \end{aligned}$$

Replacing n by $n-2k-s$, we get

$$(3.22) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{[n/2]} \sum_{s=0}^{n-2k} \frac{(A)_k(-k)_s(2A+2kI)_{n-2k-s} \left(\frac{3}{2}x\right)^{n-3s}}{k!s!(n-2k-s)!} t^n.$$

Comparing the coefficients of t^n , we get the finite series representation (3.20) for Humbert matrix polynomials. \square

Theorem 3.5. Let A , $2A$ and B be matrices in $\mathbb{C}^{N \times N}$ satisfying the spectral condition (3.2). Then the Humbert matrix polynomials have the following four ad-

ditional generating matrix functions:

$$(3.23) \quad \sum_{n=0}^{\infty} ((A)_n)^{-1} h_n^A(x) t^n = \sum_{n=0}^{\infty} \frac{(3xt)^n}{n!} \\ \times {}_1F_3\left(A+nI; \frac{A+nI}{3}, \frac{A+(n+1)I}{3}, \frac{A+(n+2)I}{3}; -\left(\frac{t}{3}\right)^3\right),$$

$$(3.24) \quad \sum_{n=0}^{\infty} (B)_n ((A)_n)^{-1} h_n^A(x) t^n = \sum_{n=0}^{\infty} \frac{(3xt)^n (B)_n}{n!} \\ \times {}_4F_3\left(\frac{B+nI}{3}, \frac{B+(n+1)I}{3}, \frac{B+(n+2)I}{3}, A+nI; \right. \\ \left. \frac{A+nI}{3}, \frac{A+(n+1)I}{3}, \frac{A+(n+2)I}{3}; -t^3\right), \quad |-t^3| < 1,$$

$$(3.25) \quad \sum_{n=0}^{\infty} [(2A)_n]^{-1} h_n^A(x) t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \sum_{r=0}^k \frac{(-k)_r \left(\frac{3}{2}xt\right)^{n+2k}}{n!k!r!2^{2k}} \\ \times \left(\left(A + \frac{1}{2}\right)_k\right)^{-1} ((2A + (n+2k)I)_r)^{-1} \left(\frac{4t}{(3x)^2}\right)^r$$

and

$$(3.26) \quad \sum_{n=0}^{\infty} (B)_n ((2A)_n)^{-1} h_n^A(x) t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \sum_{r=0}^k \frac{(-k)_r \left(\frac{3}{2}xt\right)^{n+2k}}{n!k!r!2^{2k}} (B)_{n+2k} \\ \times (B + (n+2k)I)_r \left(\left(A + \frac{1}{2}\right)_k\right)^{-1} ((2A + (n+2k)I)_r)^{-1} \left(\frac{4t}{(3x)^2}\right)^r.$$

Proof. From (3.1) we get

$$(3.27) \quad \sum_{n=0}^{\infty} ((A)_n)^{-1} h_n^A(x) t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{(n/3)} \frac{(-1)^k ((A)_n)^{-1} (A)_{n-2k} (3x)^{n-3k}}{k!(n-3k)!} t^n.$$

Using the result (1.5), we have

$$(3.28) \quad \sum_{n=0}^{\infty} ((A)_n)^{-1} h_n^A(x) t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{(-1)^k ((A)_{n+3k})^{-1} (A)_{n+k} (3x)^n}{k!n!} t^{n+3k}.$$

Using (1.3) it is easy to show that

$$(3.29) \quad (A)_{n+k} = (A)_n (A+nI)_k, \quad (A)_{n+3k} = (A)_n (A+nI)_{3k}$$

and using Gauss's multiplication theorem, we can write

$$(3.30) \quad (A + nI)_{3k} = 3^{3k} \left(\frac{A + nI}{3}\right)_k \left(\frac{A + (n+1)I}{3}\right)_k \left(\frac{A + (n+2)I}{3}\right)_k.$$

From (3.28), (3.29) and (3.30), we get

$$\begin{aligned} \sum_{n=0}^{\infty} ((A)_n)^{-1} h_n^A(x) t^n &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{1}{3^{3k} k! n!} \left((-1)^k \left(\left(\frac{A + nI}{3} \right)_k \right)^{-1} \right. \\ &\quad \times \left. \left(\left(\frac{A + (n+1)I}{3} \right)_k \right)^{-1} \left(\left(\frac{A + (n+2)I}{3} \right)_k \right)^{-1} (A + nI)_k (3x)^n \right) t^{n+3k} \\ &= \sum_{n=0}^{\infty} \frac{(3xt)^n}{n!} \sum_{k=0}^{\infty} \frac{1}{k! 3^{3k}} \left((-1)^k (A + nI)_k \left(\left(\frac{A + nI}{3} \right)_k \right)^{-1} \right. \\ &\quad \times \left. \left(\left(\frac{A + nI + I}{3} \right)_k \right)^{-1} \left(\left(\frac{A + nI + 2I}{3} \right)_k \right)^{-1} \right) t^{3k}, \end{aligned}$$

which is equivalent to (3.23).

Next, for A and $B \in \mathbb{C}^{N \times N}$ we obtain another generating matrix function for the Humbert matrix polynomials. Indeed, using (3.29) and (3.30), we find that

$$\begin{aligned} \sum_{n=0}^{\infty} (B)_n ((A)_n)^{-1} h_n^A(x) t^n &= \sum_{n=0}^{\infty} \sum_{k=0}^{(n/3)} \frac{(-1)^k (B)_n ((A)_n)^{-1} (A)_{n-2k} (3x)^{n-3k}}{k! (n-3k)!} t^n \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{(-1)^k (B)_{n+3k} ((A)_{n+3k})^{-1} (A)_{n+k} (3x)^n}{k! n!} t^{n+3k} \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{1}{k! n!} \left((-1)^k (B)_n (B + nI)_{3k} ((A)_n)^{-1} \right. \\ &\quad \times \left. ((A + nI)_{3k})^{-1} (A)_n (A + nI)_k (3x)^n \right) t^{n+3k} \\ &= \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{(-1)^k (B)_n \left(\frac{1}{3} (B + nI) \right)_k \left(\frac{1}{3} (B + (n+1)I) \right)_k \left(\frac{1}{3} (B + (n+2)I) \right)_k}{k! n!} \\ &\quad \times \left(\left(\frac{A + nI}{3} \right)_k \right)^{-1} \left(\left(\frac{A + (n+1)I}{3} \right)_k \right)^{-1} \left(\left(\frac{A + (n+2)I}{3} \right)_k \right)^{-1} \\ &\quad \times (A + nI)_k (3x)^n t^{n+3k} \\ &= \sum_{n=0}^{\infty} \frac{(B)_n (3xt)^n}{n!} {}_4F_3 \left(\frac{B + nI}{3}, \frac{B + (n+1)I}{3}, \frac{B + (n+2)I}{3}, \right. \\ &\quad \left. A + nI; \frac{A + nI}{3}, \frac{A + (n+1)I}{3}, \frac{A + (n+2)I}{3}; -t^3 \right), \end{aligned}$$

which leads to (3.25). The proof of (3.26) and (3.27) can be done similarly to that of (3.24) and (3.25) by using (3.21). \square

Expansions of Humbert matrix polynomials in series of Legendre, Gegenbauer, Hermite, Laguerre and modified Laguerre matrix polynomials relevant to the present investigation are summarized in the following theorem.

Theorem 3.6. *Let A be a matrix in $\mathbb{C}^{N \times N}$ satisfying the spectral condition (3.2). Expansions of Humbert matrix polynomials in series of Legendre, Gegenbauer, Hermite, Laguerre and modified Laguerre matrix polynomials are given by*

$$(3.31) \quad h_n^A(x) = \sum_{k=0}^{[n/3]} \sum_{r=0}^{[(n-3k)/2]} \frac{(-1)^k (A)_{n-2k} (2n - 4r - 6k + 1)}{k!r! \left(\frac{3}{2}\right)_{n-r-3k}} \times (\sqrt{2A})^{3k-n} P_{n-2r-3k}(3x, A)$$

where $P_n(x, A)$ is the Legendre matrix polynomial [35];

$$(3.32) \quad h_n^A(x) = \sum_{k=0}^{[n/3]} \sum_{r=0}^{[(n-3k)/2]} \frac{(-1)^k (A)_{n-2k}}{k!r!(n-2r-3k)!} (\sqrt{2A})^{3k-n} H_{n-2r-3k}(3x, A)$$

where $H_n(x, A)$ stands for the Hermite matrix polynomial [10], [15];

$$(3.33) \quad h_n^A(x) = \sum_{k=0}^{[n/3]} \sum_{r=0}^{[(n-3k)/2]} \frac{(-1)^k (A)_{n-2k} (A + (n-3k-2r)I)}{k!r!} \times ((A)_{n+1-3k-r})^{-1} C_{n-3k-2r}^A\left(\frac{3x}{2}\right)$$

where $C_n^A(x)$ stands for the Gegenbauer matrix polynomial [27];

$$(3.34) \quad h_n^A(x) = \sum_{k=0}^{[n/3]} \sum_{r=0}^{n-3k} \frac{1}{k!(n-r-3k)!} ((-1)^{k+r} \lambda^{-n+3k} (A)_{n-2k}) \times (A + I)_{n-3k} ((A + I)_r)^{-1} L_r^{(A, \lambda)}(3x)$$

where $L_n^{(A, \lambda)}(x)$ stands for the Laguerre matrix polynomial [20];

$$(3.35) \quad h_n^A(x) = \sum_{k=0}^{[n/3]} \sum_{r=0}^{n-3k} \frac{(-1)^k (A)_{n-2k}}{k!r!} \lambda^{-n+3k} (-A)_r f_{n-r-3k}^{(A, \lambda)}(3x)$$

where $f_n^{(A, \lambda)}(3x)$ stands for the modified Laguerre matrix polynomial (see [23], [33]).

Proof. Using the relation (3.1), we have

$$(3.36) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{[n/3]} \frac{(-1)^k (A)_{n-2k} (3x)^{n-3k}}{k!(n-3k)!} t^n.$$

From (1.5) and (3.36) we get

$$(3.37) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \frac{(-1)^k (A)_{n+k} (3x)^n}{k!n!} t^{n+3k}.$$

In [35], the expansion of $x^n I$ in a series of Legendre matrix polynomials has been given in the form

$$(3.38) \quad \frac{(3x)^n}{n!} I = (\sqrt{2A})^{-n} \sum_{r=0}^{[n/2]} \frac{(2n-4r+1)}{r! \left(\frac{3}{2}\right)_{n-r}} P_{n-2r}(3x, A),$$

hence we get

$$(3.39) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \sum_{r=0}^{[n/2]} \frac{(-1)^k (A)_{n+k} (2n-4r+1)}{k!r! \left(\frac{3}{2}\right)_{n-r}} \times (\sqrt{2A})^{-n} P_{n-2r}(3x, A)t^{n+3k}.$$

Replacing n by $n-3k$, we get

$$(3.40) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{[n/3]} \sum_{r=0}^{[(n-3k)/2]} \frac{(-1)^k (A)_{n-2k} (2n-4r-6k+1)}{k!r! \left(\frac{3}{2}\right)_{n-r-3k}} \times (\sqrt{2A})^{3k-n} P_{n-2r-3k}(3x, A)t^n.$$

Comparing the coefficients of t^n , we obtain (3.31).

In [15], the expansion of $x^n I$ in a series of Hermite matrix polynomials has been given in the form

$$(3.41) \quad (x\sqrt{2A})^n = \sum_{r=0}^{[n/2]} \frac{n!}{r!(n-2r)!} H_{n-2r}(x, A),$$

hence we get

$$(3.42) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \sum_{r=0}^{[n/2]} \frac{(-1)^k (A)_{n+k}}{k!r!(n-2r)!} (\sqrt{2A})^{-n} H_{n-2r}(3x, A)t^{n+3k}.$$

Replacing n by $n - 3k$, we get

$$(3.43) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\lfloor n/3 \rfloor} \sum_{r=0}^{\lfloor (n-3k)/2 \rfloor} \frac{(-1)^k (A)_{n-2k}}{k!r!(n-2r-3k)!} \\ \times (\sqrt{2A})^{3k-n} H_{n-2r-3k}(3x, A)t^n.$$

Comparing the coefficients of t^n , we obtain (3.32).

Using the result (1.17) in [16], [27], we get

$$(3.44) \quad (3x)^n I = n! \sum_{r=0}^{\lfloor n/2 \rfloor} \frac{(A + (n-2r)I)((A)_{n-r+1})^{-1}}{r!} C_{n-2r}^A \left(\frac{3}{2}x\right).$$

Using (3.37) and (3.44), we have

$$(3.45) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \sum_{r=0}^{\lfloor n/2 \rfloor} \frac{(-1)^k (A)_{n+k}}{k!r!} (A + (n-2r)I) \\ \times ((A)_{n-r+1})^{-1} C_{n-2r}^A \left(\frac{3}{2}x\right) t^{n+3k}.$$

Replacing n by $n - 3k$, we get

$$(3.46) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\lfloor n/3 \rfloor} \sum_{r=0}^{\lfloor (n-3k)/2 \rfloor} \frac{(-1)^k (A)_{n-2k}}{k!r!} (A + (n-2r-3k)I) \\ \times ((A)_{n-r-3k+1})^{-1} C_{n-2r-3k}^A \left(\frac{3}{2}x\right) t^n.$$

Comparing the coefficients of t^n , we obtain (3.33).

Using the results (1.13) in [20], [25], we have

$$(3.47) \quad (3x)^n I = n! \sum_{r=0}^n \frac{(-1)^r \lambda^{-n}}{(n-r)!} (A+I)_n ((A+I)_r)^{-1} L_r^{(A,\lambda)}(3x).$$

From (3.37) and (3.47) we get

$$(3.48) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \sum_{r=0}^n \frac{(-1)^{k+r} \lambda^{-n} (A)_{n+k}}{k!(n-r)!} (A+I)_n \\ \times ((A+I)_r)^{-1} L_r^{(A,\lambda)}(3x)t^{n+3k}.$$

Replacing n by $n - 3k$, we get

$$(3.49) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{[n/3]} \sum_{r=0}^{n-3k} \frac{(-1)^{k+r} \lambda^{-n+3k} (A)_{n-2k}}{k!(n-r-3k)!} (A+I)_{n-3k} \\ \times ((A+I)_r)^{-1} L_r^{(A,\lambda)}(3x)t^n.$$

Comparing the coefficients of t^n , we obtain (3.34).

Using the result (1.15) in [23], we have

$$(3.50) \quad (3\lambda x)^n I = n! \sum_{r=0}^n \frac{(-A)_r}{r!} f_{n-r}^{(A,\lambda)}(3x).$$

From (3.37) and (3.50) we have

$$(3.51) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{\infty} \sum_{r=0}^n \frac{(-1)^k (A)_{n+k}}{k!r!} \lambda^{-n} (-A)_r f_{n-r}^{(A,\lambda)}(3x)t^{n+3k}.$$

Replacing n by $n - 3k$, we get

$$(3.52) \quad \sum_{n=0}^{\infty} h_n^A(x)t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{[n/3]} \sum_{r=0}^{n-3k} \frac{(-1)^k (A)_{n-2k}}{k!r!} \lambda^{-n+3k} (-A)_r f_{n-r-3k}^{(A,\lambda)}(3x)t^n.$$

Comparing the coefficients of t^n , we obtain (3.35). Thus the proof is completed. \square

Theorem 3.7. *Let A be a matrix in $\mathbb{C}^{N \times N}$ satisfying the spectral condition (3.2). The Humbert matrix polynomials satisfy the formula*

$$(3.53) \quad \sum_{n_1+n_2+\dots+n_k=n} h_{n_1}^A(x)h_{n_2}^A(x)\dots h_{n_k}^A(x) = \sum_{s=0}^{[n/3]} \frac{(-1)^s (kA)_{n-2s} (3x)^{n-3s}}{s!(n-3s)!}$$

for $k \in \mathbb{N}$.

P r o o f. Using the power series of $(1 - 3xt + t^3)^{-kA}$ for $|3xt - t^3| < 1$ and making the necessary arrangements, we have

$$(3.54) \quad (1 - 3xt + t^3)^{-kA} = \sum_{n=0}^{\infty} \frac{(kA)_n (3x - t^2)^n}{n!} t^n \\ = \sum_{n=0}^{\infty} \sum_{s=0}^n \frac{(-1)^s (kA)_n (3x)^{n-s}}{s!(n-s)!} t^{n+2s} \\ = \sum_{n=0}^{\infty} \sum_{s=0}^{[n/3]} \frac{(-1)^s (kA)_{n-2s} (3x)^{n-3s}}{s!(n-3s)!} t^n.$$

In addition, we can write

$$(3.55) \quad (1 - 3xt + t^3)^{-kA} = \sum_{n=0}^{\infty} \left(\sum_{n_1+n_2+\dots+n_k=n} h_{n_1}^A(x) h_{n_2}^A(x) \dots h_{n_k}^A(x) \right) t^n.$$

From (3.54) and (3.55) one can see that the proof is completed. \square

Theorem 3.8. For $k \in \mathbb{N}$ and A_1, A_2, \dots, A_k being matrices in $\mathbb{C}^{N \times N}$ satisfying the spectral condition (3.2), we have the relation

$$(3.56) \quad \sum_{n_1=0}^n \dots \sum_{n_{k-1}=0}^{n-n_1-n_2-\dots-n_{k-2}} h_{n-n_1-n_2-\dots-n_{k-1}}^{A_1}(x) h_{n_1}^{A_2}(x) \dots h_{n_{k-1}}^{A_k}(x) \\ = \sum_{s=0}^{[n/3]} \frac{(-1)^s (A_1 + A_2 + \dots + A_k)_{n-2s} (3x)^{n-3s}}{s!(n-3s)!},$$

where the matrices are assumed to be commutative.

Proof. Using the power series of $(1 - 3xt + t^3)^{-(A_1+A_2+\dots+A_k)}$ for $|3xt - t^3| < 1$ and (1.5), we obtain

$$(3.57) \quad (1 - 3xt + t^3)^{-(A_1+A_2+\dots+A_k)} = \sum_{n=0}^{\infty} \frac{(A_1 + A_2 + \dots + A_k)_n (3x - t^2)^n}{n!} t^n \\ = \sum_{n=0}^{\infty} \sum_{s=0}^n \frac{(-1)^s (A_1 + A_2 + \dots + A_k)_n (3x)^{n-s}}{s!(n-s)!} t^{n+2s} \\ = \sum_{n=0}^{\infty} \sum_{s=0}^{[n/3]} \frac{(-1)^s (A_1 + A_2 + \dots + A_k)_{n-2s} (3x)^{n-3s}}{s!(n-3s)!} t^n.$$

On the other hand, we get

$$(3.58) \quad (1 - 3xt + t^3)^{-(A_1+A_2+\dots+A_k)} \\ = (1 - 3xt + t^3)^{-A_1} (1 - 3xt + t^3)^{-A_2} \dots (1 - 3xt + t^3)^{-A_k} \\ = \left(\sum_{n_1=0}^{\infty} h_{n_1}^{A_1}(x) t^{n_1} \right) \left(\sum_{n_2=0}^{\infty} h_{n_2}^{A_2}(x) t^{n_2} \right) \dots \left(\sum_{n_k=0}^{\infty} h_{n_k}^{A_k}(x) t^{n_k} \right) \\ = \sum_{n=0}^{\infty} \sum_{n_1=0}^n \dots \sum_{n_{k-1}=0}^{n-n_1-n_2-\dots-n_{k-2}} h_{n-n_1-n_2-\dots-n_{k-1}}^{A_1}(x) h_{n_1}^{A_2}(x) \dots h_{n_{k-1}}^{A_k}(x) t^n.$$

Thus the proof is completed. \square

Theorem 3.9. Let $k \in \mathbb{N}$ and let A be a matrix in $\mathbb{C}^{N \times N}$ satisfying the spectral condition (3.2). The Humbert matrix polynomials satisfy the relation

$$(3.59) \quad h_n^A(x_1 + x_2 + \dots + x_k) = \sum_{s=0}^{\lfloor n/3 \rfloor} \frac{(-1)^s (A)_{n-2s} (3(x_1 + x_2 + \dots + x_k))^{n-3s}}{s!(n-3s)!}$$

Proof. For $|3(x_1 + x_2 + \dots + x_k)t - t^3| < 1$, using (1.2), we can write

$$(3.60) \quad \begin{aligned} (1 - 3(x_1 + x_2 + \dots + x_k)t + t^3)^{-kA} &= \sum_{n=0}^{\infty} \frac{(A)_n (3(x_1 + x_2 + \dots + x_k) - t^2)^n}{n!} t^n \\ &= \sum_{n=0}^{\infty} \sum_{s=0}^n \frac{(-1)^s (A)_n (3(x_1 + x_2 + \dots + x_k))^{n-s}}{s!(n-s)!} t^{n+2s} \\ &= \sum_{n=0}^{\infty} \sum_{s=0}^{\lfloor n/3 \rfloor} \frac{(-1)^s (kA)_{n-2s} (3(x_1 + x_2 + \dots + x_k))^{n-3s}}{s!(n-3s)!} t^n. \end{aligned}$$

On the other hand, we get

$$(3.61) \quad (1 - 3(x_1 + x_1 + \dots + x_k)t + t^3)^{-A} = \sum_{n=0}^{\infty} h_n^A(x_1 + x_2 + \dots + x_k) t^n.$$

Combining (3.60) and (3.61), the proof is completed. \square

4. GENERALIZED HUMBERT MATRIX POLYNOMIALS

The purpose of this section is to introduce a new matrix polynomial representing a generalization of the Humbert matrix polynomials in (3.1). For a positive integer m , let A be a matrix in $\mathbb{C}^{N \times N}$ satisfying the spectral condition (3.2). Then we define the generalized Humbert matrix polynomials by the generating matrix function in the form

$$(4.1) \quad (1 - mxt + t^m)^{-A} = \sum_{n=0}^{\infty} h_{n,m}^A(x) t^n, \quad |mxt - t^m| < 1$$

where

$$(4.2) \quad h_{n,m}^A(x) = \sum_{k=0}^{\lfloor n/m \rfloor} \frac{(-1)^k (mx)^{n-mk}}{k!(n-mk)!} (A)_{n-(m-1)k}.$$

In a forthcoming work, we will consider the problems of a unified approach to the theory of new orthogonal matrix polynomials by the technique discussed in this paper. The notation will be used here are that implied by the following generalized Humbert matrix polynomials and generating matrix function definitions: For the generalized Humbert matrix polynomials of index two, three and p in terms of series they are represented as follows:

$$(4.3) \quad h_{n,m}^A(x,y) = \sum_{k=0}^{[n/3]} \sum_{r=0}^{[m/3]} \frac{(-1)^{k+r} (3x)^{n-3k} (3y)^{m-3r}}{k!r!(n-3k)!(m-3r)!} (A)_{n+m-2(k+r)},$$

$$\sum_{n=0}^{\infty} \sum_{m=0}^{\infty} h_{n,m}^A(x,y) t^n u^m = (1 - 3xt - 3yu + t^3 + u^3)^{-A},$$

$$|3xt + 3yu - t^3 - u^3| < 1,$$

$$(4.4) \quad h_{n,m,p}^A(x,y,z) = \sum_{k=0}^{[n/3]} \sum_{r=0}^{[m/3]} \sum_{s=0}^{[p/3]} \frac{(-1)^{k+r+s} (3x)^{n-3k} (3y)^{m-3r} (3z)^{p-3s}}{k!r!s!(n-3k)!(m-3r)!(p-3s)!}$$

$$\times (A)_{n+m+p-2(k+r+s)},$$

$$\sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sum_{p=0}^{\infty} h_{n,m,p}^A(x,y,z) t^n u^m v^p = (1 - 3xt - 3yu - 3zv + t^3 + u^3 + v^3)^{-A},$$

$$|3xt + 3yu + 3zv - t^3 - u^3 - v^3| < 1$$

and

$$(4.5) \quad h_{n_1, n_2, \dots, n_p}^A(x) = \sum_{k_i=0, i=1,2,\dots,p}^{[n_i/3]} \frac{(-1)^{\sum k_i} (3x)^{\sum n_i - 3 \sum k_i}}{\prod_{i=1}^p (k_i)! \prod_{i=1}^p (n_i - 3k_i)!} (A)_{\sum n_i - 2 \sum k_i},$$

$$\sum_{n_1=0}^{\infty} \sum_{n_2=0}^{\infty} \dots \sum_{n_p=0}^{\infty} h_{n_1, n_2, \dots, n_p}^A(x) t_1^{n_1} t_2^{n_2} \dots t_p^{n_p}$$

$$= (1 - 3x(t_1 + t_2 + \dots + t_p) + (t_1^3 + t_2^3 + \dots + t_p^3))^{-A},$$

where $|3x(t_1 + t_2 + \dots + t_p) - (t_1^3 + t_2^3 + \dots + t_p^3)| < 1$ and \sum denotes $\sum_{i=1}^p$.

Finally, we consider $P_n(m, x, y, c, A)$ which is defined by

$$(4.6) \quad (c - mxt + yt^m)^{-A} = \sum_{n=0}^{\infty} P_n(m, x, y, c, A) t^n,$$

where m is a positive integer and all parameters satisfy the condition $|mxt - yt^m| < |c|$ (see [4]). This leads us to define a new class of matrix polynomials $P_n(m, x, y, c, A)$

by the relation

$$(4.7) \quad (1 - ax + bx^l t^m)^{-A} = \sum_{n=0}^{\infty} P_n^l(m, x, a, b, A) t^n$$

where a, b, m and l are parameters satisfying the condition $|ax - bx^l t^m| < 1$.

From (4.7) we have

$$(4.8) \quad P_n^l(m, x, a, b, A) = \sum_{k=0}^{[n/m]} \frac{(-1)^k a^{n-km} b^k x^{n-(m-l)k}}{k!(n-mk)!} (A)_{n-(m-1)k}.$$

A generalization of various matrix polynomials mentioned above is provided by the following definition. This definition includes Gegenbauer, Legendre, Chebyshev, Pincherle, Kinney and Humbert matrix polynomials.

$$(4.9) \quad (c - ax + bt^m(2x - 1)^d)^{-A} = \sum_{n=0}^{\infty} P_{n,m,a,b,c,d}^A(x) t^n, \\ |ax - bt^m(2x - 1)^d| < |c|$$

where m and a are positive integers and the other parameters are unrestricted in general.

From (4.9) we get

$$(4.10) \quad P_{n,m,a,b,c,d}^A(x) = \sum_{k=0}^{[n/m]} \frac{1}{k!(n-mk)!} ((-1)^k c^{-A-nI+(m-1)kI} \\ \times (ax)^{n-(m-l)k} (b(2x-1)^d)^k) (A)_{n-(m-1)k}.$$

Setting $m = 3, c = 1$ in (4.6), we get the series representation in (3.1). Also, if we set $a = m, b = c = 1$ in (4.7), we get (4.1).

In conclusion, we remark that it would be easy to extend these properties to certain polynomials with two variables; for instance, just as with the generalized Humbert matrix polynomials, we can start from the expansion

$$(4.11) \quad (1 - 3xt - 3yu + t^3 + u^3)^{-A} = \sum_{m,n=0}^{\infty} h_{m,n}^A(x, y) t^m u^n, \\ |3xt + 3yu - t^3 - u^3| < 1$$

and obtain formulae such as

$$(4.12) \quad \frac{\partial}{\partial x} h_{m,n+1}^A(x, y) = \frac{\partial}{\partial y} h_{m+1,n}^A(x, y),$$

$$(4.13) \quad \frac{\partial}{\partial x} h_{m,n-2}^A(x, y) - x \frac{\partial}{\partial x} h_{m,n}^A(x, y) + n h_{m,n}^A(x, y) = \mathbf{0}, \quad n \geq 2$$

and

$$(4.14) \quad \frac{\partial}{\partial y} h_{m-2,n}^A(x, y) - y \frac{\partial}{\partial y} h_{m,n}^A(x, y) + m h_{m,n}^A(x, y) = \mathbf{0}, \quad n \geq 2.$$

There are many ways of investigating the generalized classes of Humbert matrix polynomials. Starting from the modified forms of the generating matrix function of Humbert matrix polynomials is one of the direct methods clearly offering some directions to develop more researches and studies in that area.

5. OPEN PROBLEM

One can use the same class of new integral representations, operational methods and the property of orthogonality for the generalized Humbert matrix polynomials, from which a variety of interesting results follows as special cases. Hence, new results and further applications can be obtained. Further results and applications will be discussed in a forthcoming work.

Acknowledgement. The author expresses his sincere appreciation to Mrs. Shimaa Ibrahim Moustafa Abdal-Rahman, (Department of Mathematics, Faculty of Science, Assiut University) for her kind interest, encouragements, help, suggestions, and comments.

The author would like to thank the referees for their valuable comments and suggestions which have led to better presentation of a paper.

References

- [1] *R. Aktaş*: A new multivariable extension of Humbert matrix polynomials. AIP Conference Proceedings 1558: Proc. Int. Conf. on Numerical Analysis and Applied Mathematics (T. Simos et al., eds.). AIP Publishing LLC, Rhodes, Greece, 2013, pp. 1128–1131. [doi](#)
- [2] *R. Aktaş*: A note on multivariable Humbert matrix polynomials. Gazi Univ. J. Sci. *27* (2014), 747–754.
- [3] *R. Aktaş, B. Çekim, A. Çevik*: Extended Jacobi matrix polynomials. Util. Math. *92* (2013), 47–64. [zbl](#) [MR](#)
- [4] *R. Aktaş, B. Çekim, R. Şahin*: The matrix version for the multivariable Humbert polynomials. Miskolc Math. Notes *13* (2012), 197–208. [zbl](#) [MR](#)
- [5] *A. Altın, B. Çekim*: Generating matrix functions for Chebyshev matrix polynomials of the second kind. Hacet. J. Math. Stat. *41* (2012), 25–32. [zbl](#) [MR](#)
- [6] *A. Altın, B. Çekim*: Some properties associated with Hermite matrix polynomials. Util. Math. *88* (2012), 171–181. [zbl](#) [MR](#)
- [7] *A. Altın, B. Çekim*: Some miscellaneous properties for Gegenbauer matrix polynomials. Util. Math. *92* (2013), 377–387. [zbl](#) [MR](#)
- [8] *B. Çekim, A. Altın, R. Aktaş*: Some relations satisfied by orthogonal matrix polynomials. Hacet. J. Math. Stat. *40* (2011), 241–253. [zbl](#) [MR](#)

- [9] *B. Çekim, A. Altın, R. Aktaş*: Some new results for Jacobi matrix polynomials. *Filomat* 27 (2013), 713–719. [MR](#) [doi](#)
- [10] *E. Defez, L. Jódar*: Some applications of the Hermite matrix polynomials series expansions. *J. Comput. Appl. Math.* 99 (1998), 105–117. [zbl](#) [MR](#) [doi](#)
- [11] *E. Defez, L. Jódar*: Chebyshev matrix polynomials and second order matrix differential equations. *Util. Math.* 61 (2002), 107–123. [zbl](#) [MR](#)
- [12] *E. Defez, L. Jódar, A. Law*: Jacobi matrix differential equation, polynomial solutions, and their properties. *Comput. Math. Appl.* 48 (2004), 789–803. [zbl](#) [MR](#) [doi](#)
- [13] *N. Dunford, J. T. Schwartz*: Linear Operators. Part I, General Theory. Pure and Applied Mathematics 7, A Wiley-Interscience Publishers, John Wiley & Sons, New York, 1958. [zbl](#) [MR](#)
- [14] *A. T. James*: Special functions of matrix and single argument in statistics. Theory and Application of Special Functions, Proc. Advanced Sem., Math. Res. Center, Madison, Wis. (R. A. Askey, ed.). Academic Press, New York, 1975, pp. 497–520. [zbl](#) [MR](#)
- [15] *L. Jódar, R. Company*: Hermite matrix polynomials and second order matrix differential equations. *Approximation Theory Appl.* 12 (1996), 20–30. [zbl](#) [MR](#)
- [16] *L. Jódar, R. Company, E. Ponsoda*: Orthogonal matrix polynomials and systems of second order differential equations. *Differ. Equ. Dyn. Syst.* 3 (1995), 269–288. [zbl](#) [MR](#)
- [17] *L. Jódar, J. C. Cortés*: On the hypergeometric matrix function. *J. Comput. Appl. Math.* 99 (1998), 205–217. [zbl](#) [MR](#) [doi](#)
- [18] *L. Jódar, J. C. Cortés*: Closed form general solution of the hypergeometric matrix differential equation. *Math. Comput. Modelling* 32 (2000), 1017–1028. [zbl](#) [MR](#) [doi](#)
- [19] *L. Jódar, E. Defez*: On Hermite matrix polynomials and Hermite matrix functions. *Approximation Theory Appl.* 14 (1998), 36–48. [zbl](#) [MR](#)
- [20] *L. Jódar, J. Sastre*: On Laguerre matrix polynomials. *Util. Math.* 53 (1998), 37–48. [zbl](#) [MR](#)
- [21] *L. Kargin, V. Kurt*: Some relations on Hermite matrix polynomials. *Math. Comput. Appl.* 18 (2013), 323–329. [MR](#)
- [22] *G. S. Khammash, A. Shehata*: On Humbert matrix polynomials. *Asian J. Current Eng. Maths.* 1 (2012), 232–240. <http://innovativejournal.in/index.php/ajcem/article/view/104/96>.
- [23] *S. Khan, N. A. Makboul Hassan*: 2-variable Laguerre matrix polynomials and Lie-algebraic techniques. *J. Phys. A, Math. Theor.* 43 (2010), Article ID 235204, 21 pages. [zbl](#) [MR](#) [doi](#)
- [24] *M. S. Metwally, M. T. Mohamed, A. Shehata*: On Hermite-Hermite matrix polynomials. *Math. Bohem.* 133 (2008), 421–434. [zbl](#) [MR](#)
- [25] *J. Sastre, L. Jódar*: On Laguerre matrix polynomial series. *Util. Math.* 71 (2006), 109–130. [zbl](#) [MR](#)
- [26] *A. Shehata*: On Tricomi and Hermite-Tricomi matrix functions of complex variable. *Commun. Math. Appl.* 2 (2011), 97–109. [zbl](#) [MR](#)
- [27] *A. Shehata*: A new extension of Gegenbauer matrix polynomials and their properties. *Bull. Int. Math. Virtual Inst.* 2 (2012), 29–42. [MR](#)
- [28] *A. Shehata*: On pseudo Legendre matrix polynomials. *Int. J. Math. Sci. Eng. Appl.* 6 (2012), 251–258. [MR](#)
- [29] *A. Shehata*: On Rainville’s matrix polynomials. *Sylwan J.* 158 (2014), 158–178.
- [30] *A. Shehata*: On Rice’s matrix polynomials. *Afr. Mat.* 25 (2014), 757–777. [zbl](#) [MR](#) [doi](#)
- [31] *A. Shehata*: New kinds of hypergeometric matrix functions. *British Journal of Mathematics and Computer Science* 5 (2015), 92–102. [doi](#)
- [32] *A. Shehata*: On a new family of the extended generalized Bessel-type matrix polynomials. *Mitteilungen Klosterneuburg J.* 65 (2015), 100–121.
- [33] *A. Shehata*: On modified Laguerre matrix polynomials. *J. Nat. Sci. Math.* 8 (2015), 153–166.

- [34] *F. Taşdelen, B. Çekim, R. Aktaş*: On a multivariable extension of Jacobi matrix polynomials. *Comput. Math. Appl.* *61* (2011), 2412–2423. [zbl](#) [MR](#) [doi](#)
- [35] *L. M. Upadhyaya, A. Shehata*: On Legendre matrix polynomials and its applications. *Int. Trans. Math. Sci. Comput.* *4* (2011), 291–310.
- [36] *L. M. Upadhyaya, A. Shehata*: A new extension of generalized Hermite matrix polynomials. *Bull. Malays. Math. Sci. Soc. (2)* *38* (2015), 165–179. [zbl](#) [doi](#)
- [37] *S. Varma, B. Çekim, F. Taşdelen Yeşildal*: On Konhauser matrix polynomials. *Ars Comb.* *100* (2011), 193–204. [zbl](#) [MR](#)
- [38] *S. Varma, F. Taşdelen*: Biorthogonal matrix polynomials related to Jacobi matrix polynomials. *Comput. Math. Appl.* *62* (2011), 3663–3668. [zbl](#) [MR](#) [doi](#)

Author's addresses: *Ayman Shehata*, Department of Mathematics, Faculty of Science, Assiut University, 71516 Assiut, Egypt, and Department of Mathematics, College of Science and Arts, Qassim University, King Abdulaziz Road, Unaizah, Qassim 10363, Kingdom of Saudi Arabia, e-mail: drshehata2006@yahoo.com.