

NOTE ON α -FILTERS IN DISTRIBUTIVE NEARLATTICES

ISMAEL CALOMINO, Tandil

Received September 7, 2017. Published online September 25, 2018.

Communicated by Radomír Halaš

Abstract. In this short paper we introduce the notion of α -filter in the class of distributive nearlattices and we prove that the α -filters of a normal distributive nearlattice are strongly connected with the filters of the distributive nearlattice of the annihilators.

Keywords: distributive nearlattice; annihilator; α -filter

MSC 2010: 06A12, 03G10, 06D50

1. INTRODUCTION AND PRELIMINARIES

A nearlattice is a join-semilattice with greatest element in which every principal filter is a bounded lattice. These structures are a natural generalization of the implication algebras studied by Abbott in [1] and the bounded distributive lattices. The nearlattices form a variety and has been studied by Cornish and Hickman in [14] and [16], and by Chajda, Halaš, Kühr and Kolařík in [8], [9], [10] and [11]. A particular class of nearlattices are the distributive nearlattices. In [6] and [7], a full duality is developed for distributive nearlattices and some applications are given, and recently in [15], the author proposes a sentential logic associated with the class of distributive nearlattices.

On the other hand, Cornish in [13] introduced the notion of α -ideal in the class of distributive lattices and characterizes Stone lattices in terms of α -ideals. These results were extended to the Hilbert algebras in [4] and [5]. We can study a dual notion of α -ideal in the class of distributive nearlattices, i.e. the concept of α -filter. The main objective of this paper is to introduce the notion of α -filter in the variety of distributive nearlattices. We see that the α -filters of a normal distributive near-

This work was supported by the CONICET under Grant PIP 112-201501-00412.

lattice \mathbf{A} are strongly connected with the filters of the distributive nearlattice $R(\mathbf{A})$ of the annihilators. This result extends those obtained by Cornish.

Let $\mathbf{A} = \langle A, \vee, 1 \rangle$ be a join-semilattice with greatest element. A filter is a subset F of A such that $1 \in F$, if $a \leq b$ and $a \in F$, then $b \in F$ and if $a, b \in F$, then $a \wedge b \in F$ whenever $a \wedge b$ exists. If X is a nonempty subset of A , the smallest filter containing X is called the filter generated by X and will be denoted by $F(X)$. A filter G is said to be finitely generated if $G = F(X)$ for some finite nonempty subset X of A . If $X = \{a\}$, then $F(\{a\}) = [a] = \{x \in A: a \leq x\}$, called the principal filter of a . We denote by $\text{Fi}(A)$ the set of all filters of A . A subset I of A is called an ideal if for every $a, b \in A$, if $a \leq b$ and $b \in I$, then $a \in I$ and for all $a, b \in I$, $a \vee b \in I$. We say that a nonempty proper ideal P is prime if for every $a, b \in A$, $a \wedge b \in P$ implies $a \in P$ or $b \in P$ whenever $a \wedge b$ exists. We denote by $\text{Id}(A)$ and $\text{X}(A)$ the set of all ideals and prime ideals of A , respectively. Finally, we say that a nonempty ideal I of A is maximal if it is proper and for every $J \in \text{Id}(A)$, if $I \subseteq J$, then $J = I$ or $J = A$. We denote by $\text{Idm}(A)$ the set of all maximal ideals of A . Note that every maximal ideal is prime.

Definition 1. Let \mathbf{A} be a join-semilattice with greatest element. Then \mathbf{A} is a *nearlattice* if each principal filter is a bounded lattice with respect to the induced order.

Note that the operation meet is defined only in a corresponding principal filter. We indicate this fact by indices, i.e. \wedge_a denotes the meet in $[a]$. Then the operation meet is not defined everywhere. However, the nearlattices can be regarded as total algebras through a ternary operation. This fact was first proved by Hickman in [16] and independently by Chajda and Kolařík in [11]. Araújo and Kinyon in [2] found a smaller equational base.

Theorem 2 ([2]). *Let \mathbf{A} be a nearlattice. Let $m: A^3 \rightarrow A$ be a ternary operation given by $m(x, y, z) = (x \vee z) \wedge_z (y \vee z)$. The following identities are satisfied:*

- (1) $m(x, y, x) = x$,
- (2) $m(m(x, y, z), m(y, m(u, x, z), z), w) = m(w, w, m(y, m(x, u, z), z))$,
- (3) $m(x, x, 1) = 1$.

Conversely, let $\mathbf{A} = \langle A, m, 1 \rangle$ be an algebra of type $(3, 0)$ satisfying the identities (1)–(3). If we define $x \vee y = m(x, x, y)$, then \mathbf{A} is a join-semilattice with greatest element. Moreover, for each $a \in A$, $[a]$ is a bounded lattice, where for every $x, y \in [a]$ their infimum is $x \wedge_a y = m(x, y, a)$. Hence, \mathbf{A} is a nearlattice.

Definition 3. Let \mathbf{A} be a nearlattice. Then \mathbf{A} is *distributive* if each principal filter is a bounded distributive lattice with respect to the induced order.

Example 4 ([1]). An implication algebra can be defined as a join-semilattice with greatest element such that each principal filter is a Boolean lattice. If $\mathbf{A} = \langle A, \rightarrow, 1 \rangle$ is an implication algebra, then the join of two elements x and y is given by $x \vee y = (x \rightarrow y) \rightarrow y$ and for each $a \in A$, $[a] = \{x \in A : a \leq x\}$ is a Boolean lattice, where for $x, y \in [a]$ the meet is given by $x \wedge_a y = (x \rightarrow (y \rightarrow a)) \rightarrow a$ and $x \rightarrow a$ is the complement of x in $[a]$. Thus, every implication algebra is a distributive nearlattice.

From the results given in [14], we have the following characterization of the filter generated by a nonempty subset X in a distributive nearlattice \mathbf{A} :

$$F(X) = \{a \in A : \exists x_1, \dots, x_n \in [X], \exists x_1 \wedge \dots \wedge x_n, a = x_1 \wedge \dots \wedge x_n\}.$$

In [3] it was shown that if \mathbf{A} is a distributive nearlattice, then the set of all filters $\text{Fi}(\mathbf{A}) = \langle \text{Fi}(A), \vee, \bar{\wedge}, \rightarrow, \{1\}, A \rangle$ is a Heyting algebra, where the least element is $\{1\}$, the greatest element is A , $G \vee H = F(G \cup H)$, $G \bar{\wedge} H = G \cap H$ and

$$(\star) \quad G \rightarrow H = \{a \in A : [a] \cap G \subseteq H\}$$

for all $G, H \in \text{Fi}(A)$. So, the pseudocomplement of $F \in \text{Fi}(A)$ is $F^* = F \rightarrow \{1\}$.

Theorem 5 ([9]). *Let \mathbf{A} be a distributive nearlattice. Let $I \in \text{Id}(A)$ and let $F \in \text{Fi}(A)$ such that $I \cap F = \emptyset$. Then there exists $P \in \text{X}(A)$ such that $I \subseteq P$ and $P \cap F = \emptyset$.*

The following definition given in [3] is an alternative definition of relative annihilator in distributive nearlattices different from that given in [10].

Definition 6. Let \mathbf{A} be a join-semilattice with greatest element and $a, b \in A$. The *annihilators of a relative to b* is the set

$$a \circ b = \{x \in A : b \leq x \vee a\}.$$

In particular, the relative annihilator $a^\top = a \circ 1 = \{x \in A : x \vee a = 1\}$ is called the *annihilator of a* .

It follows that a nearlattice \mathbf{A} is distributive if and only if $a \circ b \in \text{Fi}(A)$ for all $a, b \in A$. Also note that by (\star) , we have that $[a]^* = \{x \in A : x \vee a = 1\}$, i.e. $[a]^* = a^\top$, which is the dual notion of annulet given by Cornish in [13]. The following result will be useful.

Lemma 7 ([3]). *Let \mathbf{A} be a distributive nearlattice. Let $a, b \in A$ and $I \in \text{Id}(A)$.*

- (1) $I \cap a^\top = \emptyset$ if and only if there exists $U \in \text{Idm}(A)$ such that $I \subseteq U$ and $a \in U$.
- (2) $U \in \text{Idm}(A)$ if and only if for every $a \in A$, $a \notin U$ if and only if $U \cap a^\top \neq \emptyset$.

We are interested in a particular class of distributive nearlattices which generalize the normal lattices given in [12].

Definition 8. Let \mathbf{A} be a distributive nearlattice. Then \mathbf{A} is *normal* if each prime ideal is contained in a unique maximal ideal.

Theorem 9 ([3]). *Let \mathbf{A} be a distributive nearlattice. The following conditions are equivalent:*

- (1) \mathbf{A} is normal,
- (2) $(a \vee b)^\top = a^\top \vee b^\top$ for all $a, b \in A$.

2. α -FILTERS

In this section we study the notion of α -filter in the class of distributive nearlattices. First, we see some characteristics of annihilators. Let \mathbf{A} be a distributive nearlattice, $a \in A$ and we consider the set

$$a^{\top\top} = \{y \in A : \forall x \in a^\top, y \vee x = 1\}.$$

Lemma 10. *Let \mathbf{A} be a distributive nearlattice. The following properties are satisfied for every $a, b \in A$:*

- (1) $[a] \subseteq a^{\top\top}$.
- (2) $a^{\top\top\top} = a^\top$.
- (3) $a \leq b$ implies $a^\top \subseteq b^\top$.
- (4) $a^\top \subseteq b^\top$ if and only if $b^{\top\top} \subseteq a^{\top\top}$.
- (5) $(a \wedge b)^\top = a^\top \cap b^\top$ whenever $a \wedge b$ exists.
- (6) $(a \vee b)^{\top\top} = a^{\top\top} \cap b^{\top\top}$.

Proof. We prove only the assertions (2), (4) and (6).

(2) Let $y \in a^{\top\top\top}$. Thus, for every $x \in a^{\top\top}$ we have $y \vee x = 1$. In particular, $a \in a^{\top\top}$ and $y \vee a = 1$. Therefore $y \in a^\top$. The reciprocal is similar.

(4) Suppose that $a^\top \subseteq b^\top$. Let $y \in b^{\top\top}$. If $x \in a^\top$, then $x \in b^\top$ and $y \vee x = 1$. So, $y \in a^{\top\top}$ and $b^{\top\top} \subseteq a^{\top\top}$. Conversely, suppose that $b^{\top\top} \subseteq a^{\top\top}$ and let $x \in a^\top$. Since $b \in b^{\top\top}$, $b \in a^{\top\top}$ and $b \vee x = 1$. Therefore $x \in b^\top$ and $a^\top \subseteq b^\top$.

(6) Since $a, b \leq a \vee b$, we have $(a \vee b)^{\top\top} \subseteq a^{\top\top}, b^{\top\top}$ and $(a \vee b)^{\top\top} \subseteq a^{\top\top} \cap b^{\top\top}$. Let $y \in a^{\top\top} \cap b^{\top\top}$ and suppose that $y \notin (a \vee b)^{\top\top}$. Then there is $x \in (a \vee b)^\top$ such that $y \vee x < 1$ and by Theorem 5, there exists $P \in X(A)$ such that $y \vee x \in P$. So, $x, y \in P$. Since $y \in a^{\top\top} \cap b^{\top\top}$, we have that for every $z \in a^\top$, $y \vee z = 1$ and for every $w \in b^\top$, $y \vee w = 1$. On the other hand, as $x \in (a \vee b)^\top$, it follows that

$a \vee b \vee x = 1$ and $a \vee x \in b^\top$. Consequently, $y \vee a \vee x = 1$. We have two cases: if $P \cap a^\top \neq \emptyset$, then there is $t \in a^\top$ such that $t \in P$. Thus, $y \vee t = 1 \in P$, which is a contradiction. If $P \cap a^\top = \emptyset$, then by Lemma 7 there exists $U \in \text{Idm}(A)$ such that $P \subseteq U$ and $a \in U$. So, $x, y, a \in U$ and $y \vee a \vee x = 1 \in U$, which is a contradiction. Therefore, we conclude that $(a \vee b)^{\top\top} = a^{\top\top} \cap b^{\top\top}$. \square

If \mathbf{A} is a distributive nearlattice, then an element $a \in A$ is *dense* if $a^\top = \{1\}$. We denote by $D(A)$ the set of all dense elements of A . By Lemma 10, it is easy to prove that $D(A) \in \text{Id}(A)$ and $a^{\top\top} \in \text{Fi}(A)$ for all $a \in A$. The following result gives an equivalence of the implication algebras in terms of annihilators.

Theorem 11. *Let \mathbf{A} be a distributive nearlattice. The following conditions are equivalent:*

- (1) \mathbf{A} is an implication algebra,
- (2) $[a] \vee a^\top = A$ for all $a \in A$.

Proof. (1) \Rightarrow (2): Suppose that \mathbf{A} is an implication algebra. By the results developed in [1], we know that $X(A) = \text{Idm}(A)$. Let $a \in A$. Obviously $[a] \vee a^\top \subseteq A$. We prove the other inclusion. Let $c \in A$ and suppose that $c \notin [a] \vee a^\top$. So, by Theorem 5 there exists $P \in X(A)$ such that $c \in P$ and $P \cap ([a] \vee a^\top) = \emptyset$. Then $a \notin P$ and $P \cap a^\top = \emptyset$. Thus, P is maximal and by Lemma 7 it follows that $P \cap a^\top \neq \emptyset$, which is a contradiction. Therefore $[a] \vee a^\top = A$.

(2) \Rightarrow (1): Let $a \in A$ and $b \in [a]$ such that $b \neq a$ and $b \neq 1$. Let us prove that b has a complement in $[a]$. We know that $a \in [b] \vee b^\top = F([b] \cup b^\top)$. If only there is $x \in [b]$ such that $a = x$, then $b \leq x = a$ and $b = a$, which is a contradiction. On the other hand, if only there is $x \in b^\top$ such that $a = x$, then $x \vee b = a \vee b = 1$. Since $a \leq b$, it follows that $a \vee b = b$ and $b = 1$, which is a contradiction. Thus, there exists $x \in [b]$ and there exists $y \in b^\top$ such that $x \wedge y$ exists and $a = x \wedge y$. Then

$$a = a \wedge b = (x \wedge y) \wedge b = (x \wedge b) \wedge y = b \wedge y,$$

i.e. $a = b \wedge y$. Moreover, $y \in b^\top$ and $b \vee y = 1$. As $y \in [a]$, then y is the complement of b in $[a]$ and \mathbf{A} is an implication algebra. \square

Let \mathbf{A} be a normal distributive nearlattice and we consider the family

$$R(A) = \{a^\top : a \in A\}.$$

Let $\overline{m}: R(A)^3 \rightarrow R(A)$ be a map given by $\overline{m}(a^\top, b^\top, c^\top) = (a^\top \vee c^\top) \wedge (b^\top \vee c^\top)$. By Theorems 9 and 2 and Lemma 10, it follows that the structure

$$R(\mathbf{A}) = \langle R(A), \overline{m}, A \rangle$$

is a distributive nearlattice.

Corollary 12. Let \mathbf{A} be a normal distributive nearlattice. Then the relation θ^\top on A defined by

$$(*) \quad (a, b) \in \theta^\top \quad \text{if only if} \quad a^\top = b^\top$$

is a congruence on \mathbf{A} .

Corollary 13. Let \mathbf{A} be a normal distributive nearlattice and θ^\top be the congruence given by (*). Then $R(\mathbf{A})$ is isomorphic to \mathbf{A}/θ^\top .

Proof. Let $\varrho: A \rightarrow R(A)$ be the map defined by $\varrho(a) = a^\top$. By Theorem 9 and Lemma 10 we have that $\varrho(m(a, b, c)) = \overline{m}(\varrho(a), \varrho(b), \varrho(c))$, where the ternary operation $m(a, b, c)$ is given by Theorem 2. So, ϱ is an homomorphism onto such that $\theta^\top = \text{Ker}(\varrho)$. It follows by Isomorphism Theorem. \square

Example 14. Let \mathbf{A} be the normal distributive nearlattice from Figure 1. Then $R(A) = \{1^\top, a^\top, b^\top, c^\top\}$. On the other hand, the congruence θ^\top is given by the partition $\{1\}, \{b\}, \{a, d\}$ and $\{c, e\}$. Hence, $R(\mathbf{A})$ and \mathbf{A}/θ^\top are isomorphic.




Figure 1.

Definition 15. Let \mathbf{A} be a distributive nearlattice and $F \in \text{Fi}(A)$. We say that F is an α -filter if $a^{\top\top} \subseteq F$ for all $a \in F$.

We denote by $\text{Fi}_\alpha(A)$ the set of all α -filters of A .

Example 16. If \mathbf{A} is a normal distributive nearlattice, then $\text{Ker}(\theta^\top)$ is an α -filter.

Example 17. If \mathbf{A} is a distributive nearlattice, then a^\top is an α -filter for all $a \in A$. Let $x \in a^\top$. We prove that $x^{\top\top} \subseteq a^\top$. If $y \in x^{\top\top}$, then $x^\top \subseteq y^\top$ and since a^\top is a filter, we have $x \vee y \in a^\top$ and $x \vee y \vee a = 1$, i.e. $y \vee a \in x^\top$. So, $y \vee a \in y^\top$ and $y \vee a = 1$. It follows that $y \in a^\top$ and a^\top is an α -filter.

Remark 18. Not every filter is an α -filter. In Example 14, we consider the filter $F = \{1, a, b\}$. Thus, $a^{\top\top} = \{1, a, d\}$ and $a^{\top\top} \not\subseteq F$.

Theorem 19. Let \mathbf{A} be a distributive nearlattice and $F \in \text{Fi}(A)$. The following conditions are equivalent:

- (1) F is an α -filter.
- (2) If $a^\top = b^\top$ and $a \in F$, then $b \in F$ for all $a, b \in A$.
- (3) $F = \bigcup \{a^{\top\top} : a \in F\}$.

Proof. (1) \Rightarrow (2): Let $a, b \in A$ such that $a^\top = b^\top$ and $a \in F$. Then $a^{\top\top} = b^{\top\top}$ and since F is an α -filter, $a^{\top\top} \subseteq F$. Then $b \in b^{\top\top}$ and $b^{\top\top} \subseteq F$, i.e. $b \in F$.

(2) \Rightarrow (3): Since $a \in a^{\top\top}$ for all $a \in A$, we have $F \subseteq \bigcup \{a^{\top\top} : a \in F\}$. We see the other inclusion. If $x \in \bigcup \{a^{\top\top} : a \in F\}$, then there is $b \in F$ such that $x \in b^{\top\top}$. So, $b^\top \subseteq x^\top$ and $x^{\top\top} \subseteq b^{\top\top}$. Then by Lemma 10, $x^{\top\top} = x^{\top\top} \cap b^{\top\top} = (x \vee b)^{\top\top}$ and $x^\top = (x \vee b)^\top$. As $x \vee b \in F$, by hypothesis we have $x \in F$.

(3) \Rightarrow (1): Let $b \in F$. If $x \in b^{\top\top}$, then $x \in \bigcup \{a^{\top\top} : a \in F\}$ and $x \in F$. Therefore $b^{\top\top} \subseteq F$ and F is an α -filter. \square

Theorem 20. Let \mathbf{A} be a normal distributive nearlattice and $F \in \text{Fi}(A)$. Then

$$\alpha(F) = \{x \in A : \exists a \in F, a^\top \subseteq x^\top\}$$

is the smallest α -filter containing F .

Proof. It is clear that $F \subseteq \alpha(F)$. Let $x, y \in A$ such that $x \leq y$ and $x \in \alpha(F)$. Then by Lemma 10, $x^\top \subseteq y^\top$ and there exists $a \in F$ such that $a^\top \subseteq x^\top$. So, $a^\top \subseteq y^\top$ and $y \in \alpha(F)$. Let $x, y \in \alpha(F)$ and suppose that $x \wedge y$ exists. Then there exist $a, b \in F$ such that $a^\top \subseteq x^\top$ and $b^\top \subseteq y^\top$. Since F is a filter, $m(a, b, x \wedge y) \in F$, where the ternary operation $m(a, b, x \wedge y)$ is given by Theorem 2. On the other hand, $a^\top \vee (x \wedge y)^\top \subseteq x^\top$ and $b^\top \vee (x \wedge y)^\top \subseteq y^\top$. As \mathbf{A} is normal,

$$m(a, b, x \wedge y)^\top = \overline{m(a^\top, b^\top, (x \wedge y)^\top)} \subseteq x^\top \bar{\wedge} y^\top = (x \wedge y)^\top.$$

Thus, $m(a, b, x \wedge y)^\top \subseteq (x \wedge y)^\top$ and $x \wedge y \in \alpha(F)$. Then $\alpha(F)$ is a filter. Let $x \in \alpha(F)$. We see that $x^{\top\top} \subseteq \alpha(F)$. If $y \in x^{\top\top}$, then $x^\top \subseteq y^\top$. Since $x \in \alpha(F)$, there exists $a \in F$ such that $a^\top \subseteq x^\top$. So, $a^\top \subseteq y^\top$ and $y \in \alpha(F)$. Then $x^{\top\top} \subseteq \alpha(F)$ and $\alpha(F)$ is an α -filter. Let $H \in \text{Fi}_\alpha(A)$ such that $F \subseteq H$. If $x \in \alpha(F)$, then there exists $a \in F$ such that $a^\top \subseteq x^\top$, i.e. $x^{\top\top} \subseteq a^{\top\top}$. As $a \in H$ and H is an α -filter, we have $a^{\top\top} \subseteq H$. Consequently, $x \in H$ and $\alpha(F) \subseteq H$. \square

Remark 21. Let \mathbf{A} be a normal distributive nearlattice.

- (1) Note that the map $\alpha: \text{Fi}(A) \rightarrow \text{Fi}(A)$ of Theorem 20 is a closure operator and the α -filters are closed elements with respect to α .
- (2) A proper α -filter contains non-dense elements. Indeed, if F is a proper α -filter and $x \in F \cap D(A)$, then $F = \alpha(F)$ and $x^\top = \{1\}$. Thus, there exists $a \in F$ such that $a^\top \subseteq x^\top$. So, $a^\top = \{1\}$ and $a^{\top\top} = A$. On the other hand, since F is an α -filter, $a^{\top\top} \subseteq F$, i.e. $A = F$ which is a contradiction.

Now, we define the operations of infimum $\overline{\sqcup}$, supremum $\underline{\sqcup}$, and implication \Rightarrow in $\text{Fi}_\alpha(A)$ as:

$$F \overline{\sqcup} G = F \cap G, \quad F \underline{\sqcup} G = \alpha(F \vee G), \quad F \Rightarrow G = \alpha(F \rightarrow G)$$

for each pair $F, G \in \text{Fi}_\alpha(A)$. By Theorem 20, we have that $F \overline{\sqcup} G, F \underline{\sqcup} G, F \Rightarrow G \in \text{Fi}_\alpha(A)$ for all $F, G \in \text{Fi}_\alpha(A)$. Consider the structure

$$\text{Fi}_\alpha(\mathbf{A}) = \langle \text{Fi}_\alpha(A), \underline{\sqcup}, \overline{\sqcup}, \Rightarrow, \{1\}, A \rangle.$$

Theorem 22. *Let \mathbf{A} be a normal distributive nearlattice. Then $\text{Fi}_\alpha(\mathbf{A})$ is a Heyting algebra.*

Proof. It is easy to verify that $\langle \text{Fi}_\alpha(A), \underline{\sqcup}, \overline{\sqcup}, \{1\}, A \rangle$ is a bounded lattice. Let $F, H, K \in \text{Fi}_\alpha(A)$. Suppose that $F \overline{\sqcup} H \subseteq K$. If $x \in F$, then $[x] \cap H \subseteq F \overline{\sqcup} H \subseteq K$. Thus, $[x] \cap H \subseteq K$ and $x \in H \rightarrow K$. Hence, $x \in H \Rightarrow K$ and $F \subseteq H \Rightarrow K$.

Reciprocally, we assume that $F \subseteq H \Rightarrow K$. Let $x \in F \overline{\sqcup} H$. So, $x \in F \subseteq H \Rightarrow K$ and there exists $a \in H \rightarrow K$ such that $a^\top \subseteq x^\top$. It follows that $x \vee a \in [a] \cap H \subseteq K$ and $x^\top = x^\top \vee a^\top = (x \vee a)^\top$, i.e. $x^\top = (x \vee a)^\top$ and $x \vee a \in K$. By Theorem 19, we have $x \in K$. Therefore, $F \overline{\sqcup} H \subseteq K$ and $\text{Fi}_\alpha(\mathbf{A})$ is a Heyting algebra. \square

Let \mathbf{A} be a nearlattice. Following the results developed in [15], we introduce the next notation. For each natural number n we define inductively for every $a_1, \dots, a_n, b \in A$, the element $m^{n-1}(a_1, \dots, a_n, b)$ as follows:

- (1) $m^0(a_1, b) = m(a_1, a_1, b)$,
- (2) for $n > 1$, $m^{n-1}(a_1, \dots, a_n, b) = m(m^{n-2}(a_1, \dots, a_{n-1}, b), a_n, b)$.

Then $m^{n-1}(a_1, \dots, a_n, b) = (a_1 \vee b) \wedge_b \dots \wedge_b (a_n \vee b)$ and in particular, $m^0(a_1, b) = a_1 \vee b$ and $m^1(a_1, a_2, b) = m(a_1, a_2, b)$, where the operation $m(a_1, a_2, b)$ is given by Theorem 2. We are able to formulate our main result.

Theorem 23. *Let \mathbf{A} be a normal distributive nearlattice. Then $\text{Fi}_\alpha(\mathbf{A})$ is isomorphic to the Heyting algebra $\text{Fi}(\mathbf{R}(\mathbf{A}))$.*

Proof. We consider the map $\psi: \text{Fi}_\alpha(A) \rightarrow \text{Fi}(\text{R}(A))$ defined by

$$\psi(F) = \{a^\top : a \in F\}.$$

We prove that ψ is well-defined. Let $F \in \text{Fi}_\alpha(A)$. It is clear that $1^\top \in \psi(F)$. Let $a^\top, b^\top \in \text{R}(A)$ such that $a^\top \subseteq b^\top$ and $a^\top \in \psi(F)$. Then $b^{\top\top} \subseteq a^{\top\top}$ and $a \in F$. Thus, $b \in a^{\top\top}$ and as F is an α -filter, $a^{\top\top} \subseteq F$. So, $b \in F$ and $b^\top \in \psi(F)$. Let $a^\top, b^\top \in \psi(F)$ and suppose that $a^\top \bar{\wedge} b^\top$ exists in $\text{R}(A)$, i.e. there is $c \in A$ such that $a^\top \bar{\wedge} b^\top = c^\top$. Then $a, b \in F$ and as F is a filter, $m(a, b, c) \in F$. It follows that

$$m(a, b, c)^\top = \bar{m}(a^\top, b^\top, c^\top) = (a^\top \bar{\wedge} b^\top) \vee c^\top = c^\top$$

and $c^\top \in \psi(F)$. Thus, $a^\top \bar{\wedge} b^\top \in \psi(F)$ and $\psi(F) \in \text{Fi}(\text{R}(A))$.

Let $F, H \in \text{Fi}_\alpha(A)$. It is immediate that $\psi(F \bar{\cap} H) = \psi(F) \bar{\cap} \psi(H)$. We see that $\psi(F \sqcup H) = \psi(F) \vee \psi(H)$. Let $x^\top \in \psi(F \sqcup H)$. Then $x \in \alpha(F \vee H)$ and there exists $a \in F \vee H$ such that $a^\top \subseteq x^\top$. So, there exist $x_1, \dots, x_n \in F \cup H$ such that $x_1 \wedge \dots \wedge x_n$ exists and $a = x_1 \wedge \dots \wedge x_n$. Then $x_1^\top, \dots, x_n^\top \in \psi(F) \cup \psi(H)$. On the other hand, $a^\top = (x_1 \wedge \dots \wedge x_n)^\top = x_1^\top \bar{\wedge} \dots \bar{\wedge} x_n^\top$ and $a^\top \in \psi(F) \vee \psi(H)$. Since $\psi(F) \vee \psi(H)$ is a filter, we have $x^\top \in \psi(F) \vee \psi(H)$ and $\psi(F \sqcup H) \subseteq \psi(F) \vee \psi(H)$. Conversely, if $x^\top \in \psi(F) \vee \psi(H)$, then there exist $x_1^\top, \dots, x_n^\top \in \psi(F) \cup \psi(H)$ such that $x_1^\top \bar{\wedge} \dots \bar{\wedge} x_n^\top$ exists and $x^\top = x_1^\top \bar{\wedge} \dots \bar{\wedge} x_n^\top$. It follows that $x_1, \dots, x_n \in F \cup H$ and $m^{n-1}(x_1, \dots, x_n, x) \in F \vee H$. So,

$$m^{n-1}(x_1, \dots, x_n, x)^\top = \bar{m}^{n-1}(x_1^\top, \dots, x_n^\top, x^\top) = (x_1^\top \bar{\wedge} \dots \bar{\wedge} x_n^\top) \vee x^\top = x^\top$$

and $m^{n-1}(x_1, \dots, x_n, x)^\top \subseteq x^\top$. Thus, $x \in \alpha(F \vee H) = F \sqcup H$, i.e. $x^\top \in \psi(F \sqcup H)$ and $\psi(F) \vee \psi(H) \subseteq \psi(F \sqcup H)$. Therefore, $\psi(F \sqcup H) = \psi(F) \vee \psi(H)$.

Now, we prove that $\psi(F \Rightarrow H) = \psi(F) \rightarrow \psi(H)$. Let $x^\top \in \psi(F \Rightarrow H)$. Then $x \in F \Rightarrow H = \alpha(F \rightarrow H)$ and there exists $a \in F \rightarrow H$ such that $a^\top \subseteq x^\top$. So, $[a] \cap F \subseteq H$. We see that $x^\top \in \psi(F) \rightarrow \psi(H)$, i.e. $[x^\top] \cap \psi(F) \subseteq \psi(H)$. If $y^\top \in [x^\top] \cap \psi(F)$, then $x^\top \subseteq y^\top$ and $y \in F$. Thus, $a \vee y \in [a] \cap F$ and $a \vee y \in H$. On the other hand, since $a^\top \subseteq y^\top$, we have $y^\top = (a \vee y)^\top$. As $a \vee y \in H$ and H is an α -filter, by Theorem 19, $y \in H$. Then $y^\top \in \psi(H)$ and $x^\top \in \psi(F) \rightarrow \psi(H)$. So, $\psi(F \Rightarrow H) \subseteq \psi(F) \rightarrow \psi(H)$. We prove the other inclusion. Let $x^\top \in \psi(F) \rightarrow \psi(H)$, i.e. $[x^\top] \cap \psi(F) \subseteq \psi(H)$. Then $x^\top \in \psi(F \Rightarrow H)$ if and only if $x \in \alpha(F \rightarrow H)$ if and only if there exists $a \in F \rightarrow H$ such that $a^\top \subseteq x^\top$. We see that $x \in F \rightarrow H$. If $y \in [x] \cap F$, then by Lemma 10, $x^\top \subseteq y^\top$ and $y \in F$, i.e. $y^\top \in [x^\top] \cap \psi(F)$. Since $[x^\top] \cap \psi(F) \subseteq \psi(H)$, we have $y^\top \in \psi(H)$ and $y \in H$. Then $[x] \cap F \subseteq H$ and $x \in F \rightarrow H$. Thus, $x^\top \in \psi(F \Rightarrow H)$ and $\psi(F \Rightarrow H) = \psi(F) \rightarrow \psi(H)$.

Let $\pi: \text{Fi}(\text{R}(A)) \rightarrow \text{Fi}_\alpha(A)$ be the map given by $\pi(G) = \{a: a^\top \in G\}$. By Lemma 10, it follows that $\pi(G) \in \text{Fi}_\alpha(A)$. So, ψ and π are the inverses of each other and ψ is 1-1 and onto. Therefore ψ is an isomorphism. \square

References

- [1] *J. C. Abbott*: Semi-boolean algebra. *Mat. Vesn.*, N. Ser. 4 (1967), 177–198. [zbl](#) [MR](#)
- [2] *J. Araújo, M. Kinyon*: Independent axiom systems for nearlattices. *Czech. Math. J.* 61 (2011), 975–992. [zbl](#) [MR](#) [doi](#)
- [3] *I. Calomino, S. Celani*: A note on annihilators in distributive nearlattices. *Miskolc Math. Notes* 16 (2015), 65–78. [zbl](#) [MR](#)
- [4] *S. Celani*: α -ideals and α -deductive systems in bounded Hilbert algebras. *J. Mult.-Valued Logic and Soft Computing* 21 (2013), 493–510. [zbl](#) [MR](#)
- [5] *S. Celani*: Notes on bounded Hilbert algebras with supremum. *Acta Sci. Math.* 80 (2014), 3–19. [zbl](#) [MR](#) [doi](#)
- [6] *S. Celani, I. Calomino*: Stone style duality for distributive nearlattices. *Algebra Univers.* 71 (2014), 127–153. [zbl](#) [MR](#) [doi](#)
- [7] *S. Celani, I. Calomino*: On homomorphic images and the free distributive lattice extension of a distributive nearlattice. *Rep. Math. Logic* 51 (2016), 57–73. [zbl](#) [MR](#) [doi](#)
- [8] *I. Chajda, R. Halaš*: An example of a congruence distributive variety having no near-unanimity term. *Acta Univ. M. Belii, Ser. Math.* 13 (2006), 29–31. [zbl](#) [MR](#)
- [9] *I. Chajda, R. Halaš, J. Kühr*: Semilattice Structures. *Research and Exposition in Mathematics* 30. Heldermann, Lemgo, 2007. [zbl](#) [MR](#)
- [10] *I. Chajda, M. Kolařík*: Ideals, congruences and annihilators on nearlattices. *Acta Univ. Palacki. Olomuc., Fac. Rerum Nat., Math.* 46 (2007), 25–33. [zbl](#) [MR](#)
- [11] *I. Chajda, M. Kolařík*: Nearlattices. *Discrete Math.* 308 (2008), 4906–4913. [zbl](#) [MR](#) [doi](#)
- [12] *W. H. Cornish*: Normal lattices. *J. Aust. Math. Soc.* 14 (1972), 200–215. [zbl](#) [MR](#) [doi](#)
- [13] *W. H. Cornish*: Annulets and α -ideals in distributive lattices. *J. Aust. Math. Soc.* 15 (1973), 70–77. [zbl](#) [MR](#) [doi](#)
- [14] *W. H. Cornish, R. C. Hickman*: Weakly distributive semilattices. *Acta Math. Acad. Sci. Hung.* 32 (1978), 5–16. [zbl](#) [MR](#) [doi](#)
- [15] *L. J. González*: The logic of distributive nearlattices. *Soft Computing* 22 (2018), 2797–2807. [doi](#)
- [16] *R. Hickman*: Join algebras. *Commun. Algebra* 8 (1980), 1653–1685. [zbl](#) [MR](#) [doi](#)

Author's address: Ismael Calomino, Comisión de Investigaciones Científicas—Departamento de Matemáticas, Facultad de Ciencias Exactas, Universidad Nacional del Centro, Pinto 399, 7000 Tandil, Argentina, e-mail: calomino@exa.unicen.edu.ar.