

A POWER OF A MEROMORPHIC FUNCTION SHARING  
TWO SMALL FUNCTIONS WITH A DERIVATIVE OF THE POWER

LAHIRI INDRAJIT, Kalyani, SUJOY MAJUMDER, Raiganj

Received July 8, 2020. Published online November 22, 2021.

Communicated by Grigore Sălăgean

*Abstract.* In connection to a conjecture of W. Lü, Q. Li and C. Yang (2014), we prove a result on small function sharing by a power of a meromorphic function with few poles with a derivative of the power. Our results improve a number of known results.

*Keywords:* meromorphic function; derivative; small function

*MSC 2020:* 30D35

## 1. INTRODUCTION DEFINITIONS AND RESULTS

In this paper a meromorphic function means a function that is meromorphic in the open complex plane  $\mathbb{C}$ . We use the standard notations of Nevanlinna theory; e.g.,  $N(r, f)$ ,  $m(r, f)$ ,  $T(r, f)$ ,  $N(r, a; f)$ ,  $\overline{N}(r, a; f)$ ,  $m(r, a; f)$  etc., see [7]. We denote by  $S(r, f)$  a quantity, not necessarily the same at each of its occurrences, that satisfies the condition  $S(r, f) = o\{T(r, f)\}$  as  $r \rightarrow \infty$  except possibly a set of finite linear measure.

A meromorphic function  $a = a(z)$  is called a small function of a meromorphic function  $f$  if  $T(r, a) = S(r, f)$ . Let us denote by  $S(f)$  the class of all small functions of  $f$ . Clearly  $\mathbb{C} \subset S(f)$  and if  $f$  is a transcendental function then every polynomial is a member of  $S(f)$ .

Let  $f$  and  $g$  be two non-constant meromorphic functions and  $a \in S(f) \cap S(g)$ . If  $f - a$  and  $g - a$  have the same zeros with the same multiplicities, then we say that  $f$  and  $g$  share the small function  $a$  CM (counting multiplicities) and if we do not consider the multiplicities, then we say that  $f$  and  $g$  share the small function  $a$  IM (ignoring multiplicities).

Let  $k$  be a positive integer and  $a \in S(f)$ . We use  $N_k(r, a; f)$  to denote the counting function of zeros of  $f - a$  with multiplicity not greater than  $k$ ,  $N_{(k+1)}(r, a; f)$  to denote the counting function of zeros of  $f - a$  with multiplicity greater than  $k$ . Similarly we use  $\overline{N}_k(r, a; f)$  and  $\overline{N}_{(k+1)}(r, a; f)$  to denote their respective reduced functions.

In 1996, Brück studied the relation between  $f$  and  $f'$  if an entire function  $f$  shares only one finite value CM with its derivative  $f'$  (see [1]). In this direction an interesting conjecture was proposed by Brück (see [1]), which is still open in its full generality.

**Conjecture A.** *Let  $f$  be a non-constant entire function. Suppose*

$$\varrho_1(f) := \limsup_{r \rightarrow \infty} \frac{\log \log T(r, f)}{\log r},$$

*the hyper-order of  $f$ , is not a positive integer or infinity. If  $f$  and  $f'$  share a finite value  $a$  CM, then*

$$(1.1) \quad \frac{f' - a}{f - a} = c$$

*for some nonzero constant  $c$ .*

The conjecture for the special cases

- (1)  $a = 0$ ,
- (2)  $N(r, 0; f') = S(r, f)$

had been established by Brück, see [1]. From the differential equations

$$(1.2) \quad \frac{f' - a}{f - a} = e^{z^n}, \quad \frac{f' - a}{f - a} = e^{e^z},$$

we see that when  $\varrho_1(f)$  is a positive integer or infinity the conjecture does not hold.

The conjecture for the case where  $f$  is of finite order had been proved by Gundersen and Yang (see [6]), and the case where  $f$  is of infinite order with  $\varrho_1(f) < \frac{1}{2}$  had been proved by Chen and Shon (see [3]). Recently Cao in [2] proved that the Brück conjecture is also true when  $f$  is of infinite order with  $\varrho_1(f) = \frac{1}{2}$ . But the case  $\varrho_1(f) > \frac{1}{2}$  is still open. However, the corresponding conjecture for meromorphic functions fails in general (see [6]). For example, if

$$f(z) = \frac{2e^z + z + 1}{e^z + 1},$$

then  $f$  and  $f'$  share 1 CM, but (1.1) does not hold.

It is interesting to ask what happens if  $f$  is replaced by a power of itself, say,  $f^n$  in Brück's conjecture. From (1.2) we see that the conjecture does not hold without any restriction on the hyper-order when  $n = 1$ . So we only need to focus on the problem when  $n \geq 2$ .

Perhaps Yang and Zhang (see [15]) were the first to consider the uniqueness of a power of an entire function  $F = f^n$  and its derivative  $F'$  when they share a certain value and that leads to a specific form of the function  $f$ .

Yang and Zhang in [15] proved that the Brück conjecture holds for the function  $f^n$  and the order restriction on  $f$  is not needed if  $n$  is relatively large. In fact, they proved the following result.

**Theorem A** ([15]). *Let  $f$  be a non-constant entire function,  $n (\geq 7)$  be an integer and let  $F = f^n$ . If  $F$  and  $F'$  share 1 CM, then  $F \equiv F'$ , and  $f$  assumes the form  $f(z) = ce^{z/n}$ , where  $c$  is a nonzero constant.*

Improving all the results obtained in [15], Zhang in [16] proved the following theorem.

**Theorem B** ([16]). *Let  $f$  be a non-constant entire function, and let  $n, k$  be positive integers and  $a (\neq 0, \infty)$  be a meromorphic small function of  $f$ . If  $f^n - a$  and  $(f^n)^{(k)} - a$  share 0 CM and  $n \geq k + 5$ , then  $f^n \equiv (f^n)^{(k)}$ , and  $f$  assumes the form  $f(z) = ce^{z/\lambda^n}$ , where  $c$  is a nonzero constant and  $\lambda^k = 1$ .*

In 2009, Zhang and Yang (see [17]) further improved the above result in the following manner.

**Theorem C** ([17]). *Let  $f$  be a non-constant entire function, and let  $n, k$  be positive integers and  $a (\neq 0, \infty)$  be a meromorphic small function of  $f$ . Suppose  $f^n - a$  and  $(f^n)^{(k)} - a$  share 0 CM and  $n \geq k + 2$ . Then conclusion of Theorem B holds.*

In 2010, Zhang and Yang (see [18]) further improved the above result in the following manner.

**Theorem D** ([18]). *Let  $f$  be a non-constant entire function, and let  $n, k$  be positive integers. Suppose  $f^n$  and  $(f^n)^{(k)}$  share 1 CM and  $n \geq k + 1$ . Then conclusion of Theorem B holds.*

In 2011, Lü and Yi (see [11]) proved the following extension of Theorem D.

**Theorem E** ([11]). *Let  $f$  be a transcendental entire function, and let  $n, k$  be two integers with  $n \geq k + 1$ , and let  $F = f^n$  and  $Q \neq 0$  be polynomials. If  $F - Q$  and  $F^{(k)} - Q$  share 0 CM, then  $F \equiv F^{(k)}$  and  $f(z) = ce^{wz/n}$ , where  $c$  and  $w$  are nonzero constants such that  $w^k = 1$ .*

**Remark 1.1.** It is easy to see that the condition  $n \geq k + 1$  in Theorem E is sharp by the following example.

**Example 1.1.** Let  $f(z) = e^{e^z} \int_0^z e^{-e^t} (1 - e^t) t dt$  and  $n = 1, k = 1$ . Then

$$\frac{f'(z) - z}{f(z) - z} = e^z$$

and  $f'(z) - z$  and  $f(z) - z$  share 0 CM, but  $f' \neq f$ .

In [12], Lü, Li and Yang asked the question of considering two shared polynomials in Theorem E instead of a single shared polynomial. They answered the question for the first derivative of the power of a transcendental entire function and further proposed the following conjecture:

**Conjecture B.** *Let  $f$  be a transcendental entire function, and let  $n$  be a positive integer. If  $f^n - Q_1$  and  $(f^n)^{(k)} - Q_2$  share 0 CM and  $n \geq k + 1$ , then  $(f^n)^{(k)} = f^n Q_2 / Q_1$ , where  $Q_1$  and  $Q_2$  are polynomials with  $Q_1 Q_2 \neq 0$ . If, further,  $Q_1 \equiv Q_2$ , then  $f = ce^{\omega z/n}$ , where  $c$  and  $\omega$  are nonzero constants such that  $\omega^k = 1$ .*

Recently the second author (see [13]) fully resolved Conjecture B; thus giving rise to a further investigation of the possibility of replacing in Conjecture B the shared polynomials by shared small functions. Here we on one hand solve this problem and also on the other hand we try to relax the nature of sharing of small functions, thereby improving a number of known results including that in [13].

Extending the idea of weighted sharing (see [8], [9]), Lin and Lin in [10] introduced the notion of weakly weighted sharing which is defined as follows.

**Definition 1.1** ([10]). Let  $f$  and  $g$  be two non-constant meromorphic functions sharing a “IM”, for  $a \in S(f) \cap S(g)$ , and let  $k$  be a positive integer or  $\infty$ .

- (i)  $\overline{N}_k^E(r, a)$  denotes the counting function of those zeros of  $f - a$  whose multiplicities are equal to the corresponding zeros of  $g - a$ , where both of their multiplicities are not greater than  $k$ , and where each zero is counted only once.
- (ii)  $\overline{N}_{(k)}^0(r, a)$  denotes the counting function of those zeros of  $f - a$  which are zeros of  $g - a$ , where both of their multiplicities are not less than  $k$ , and where each zero is counted only once.

**Definition 1.2** ([10]). For  $a \in S(f) \cap S(g)$ , if  $k$  is a positive integer or  $\infty$  and

$$\begin{aligned} \overline{N}_k(r, a; f) - \overline{N}_k^E(r, a) &= S(r, f), & \overline{N}_k(r, a; g) - \overline{N}_k^E(r, a) &= S(r, g); \\ \overline{N}_{(k+1)}(r, a; f) - \overline{N}_{(k+1)}^0(r, a) &= S(r, f), & \overline{N}_{(k+1)}(r, a; g) - \overline{N}_{(k+1)}^0(r, a) &= S(r, g); \end{aligned}$$

or if  $k = 0$  and

$$\overline{N}(r, a; f) - \overline{N}_0(r, a) = S(r, f), \quad \overline{N}(r, a; g) - \overline{N}_0(r, a) = S(r, g),$$

then we say  $f$  and  $g$  weakly share  $a$  with weight  $k$ . Here we write  $f, g$  share “ $(a, k)$ ” to mean that  $f, g$  weakly share  $a$  with weight  $k$ .

Obviously, if  $f$  and  $g$  share “ $(a, k)$ ”, then  $f$  and  $g$  share “ $(a, p)$ ” for any  $p$ ,  $0 \leq p \leq k$ . Also we note that  $f$  and  $g$  share a “IM” or “CM” if and only if  $f$  and  $g$  share “ $(a, 0)$ ” or “ $(a, \infty)$ ”, respectively (for the definitions of “IM” and “CM” see pages 225–226 in [14]).

We note that a rational function  $f$  with  $\overline{N}(r, \infty; f) = S(r, f)$  must be a polynomial. Also a small function of a polynomial must be a constant. Since  $k \geq 1$ , clearly if  $f$  is a polynomial, then the relation  $(f^n)^{(k)} = cf^n$  does not hold for any nonzero constant  $c$  and  $n \geq k$ . Therefore in the following theorems we assume  $f$  to be transcendental.

**Theorem 1.1.** *Let  $f$  be a transcendental meromorphic function such that  $N(r, \infty; f) = S(r, f)$  and let  $a_i = a_i(z) (\not\equiv 0, \infty)$  be small functions of  $f$ , where  $i = 1, 2$ . Let  $n$  and  $k$  be two positive integers such that  $n \geq k + 1$ . If  $f^n - a_1$  and  $(f^n)^{(k)} - a_2$  share “ $(0, 1)$ ”, then  $(f^n)^{(k)} \equiv f^n a_2 a_1^{-1}$ . Furthermore, if  $a_1 \equiv a_2$ , then  $f(z) = ce^{z\lambda/n}$ , where  $c$  and  $\lambda$  are nonzero constants such that  $\lambda^k = 1$ .*

**Theorem 1.2.** *Let  $f$  be a transcendental meromorphic function such that  $\overline{N}(r, \infty; f) = S(r, f)$  and let  $a_i = a_i(z) (\not\equiv 0, \infty)$  be small functions of  $f$ , where  $i = 1, 2$ . Let  $n$  and  $k$  be two positive integers such that  $n \geq k$ . If  $f^n - a_1$  and  $(f^n)^{(k)} - a_2$  share “ $(0, 0)$ ” and  $\overline{N}_2(r, 0; f) = S(r, f)$ , then  $(f^n)^{(k)} \equiv a_2 a_1^{-1} f^n$ . Furthermore, if  $a_1 \equiv a_2$ , then  $f^n \equiv (f^n)^{(k)}$  and  $f$  assumes the form  $f(z) = ce^{z\lambda/n}$ , where  $c$  is a nonzero constant and  $\lambda^k = 1$ .*

**Note 1.1.** If  $k \geq 2$ , then in Theorem 1.2 instead of  $\overline{N}_2(r, 0; f) = S(r, f)$  we can assume  $N_1(r, 0; f) = S(r, f)$ .

**Remark 1.2.** It is easy to see that the condition  $n \geq k + 1$  in Theorem 1.1 is sharp by the following examples.

**Example 1.2.** Let  $f(z) = e^{2z} + z$ . Then  $f - a_1$  and  $f' - a_2$  share 0 CM and  $N(r, \infty; f) = 0$ , but  $f' \not\equiv a_2 a_1^{-1} f$ , where  $a_1(z) = z + 1$  and  $a_2(z) = 3$ .

**Example 1.3.** Let  $f(z) = e^{2z} + z^2 + z$ . Then  $f - a_1$  and  $f' - a_2$  share 0 CM and  $N(r, \infty; f) = 0$ , but  $f' \not\equiv a_2 a_1^{-1} f$ , where  $a_1(z) = z^2 + z + 1$  and  $a_2(z) = 2z + 3$ .

**Example 1.4.** Let

$$f(z) = e^{e^{z^2}} + 1, \quad a_1(z) = \frac{1}{1 + e^{-z^2}}, \quad a_2(z) = -\frac{2z}{1 + e^{-z^2}}.$$

We note that

$$f(z) - a_1(z) = \frac{1}{e^{z^2} + 1} ((e^{z^2} + 1)e^{e^{z^2}} + 1)$$

and

$$f'(z) - a_2(z) = \frac{2z}{1 + e^{-z^2}} ((e^{z^2} + 1)e^{e^{z^2}} + 1).$$

Then  $f - a_1$  and  $f' - a_2$  share “ $(0, \infty)$ ” and  $N(r, \infty; f) = 0$ , but  $f \not\equiv a_2 a_1^{-1} f'$ .

Example 1.5. Let

$$f(z) = 1 - 5(z + 1) + ze^z$$

and  $a_1(z) = a_2(z) = -(4 + 4z + 5z^2)$ . We note that

$$f(z) - a_1(z) = z(e^z + 5z - 1)$$

and

$$f'(z) - a_2(z) = (z + 1)(e^z + 5z - 1).$$

Then  $f - a_1$  and  $f' - a_2$  share “ $(0, \infty)$ ” and  $N(r, \infty; f) = 0$ , but  $f \not\equiv f'$ .

Remark 1.3. It is easy to see that the conditions  $\overline{N}_2(r, 0; f) = S(r, f)$  and  $\overline{N}(r, \infty; f) = S(r, f)$  in Theorem 1.2 are essential by the following examples.

Example 1.6. Let

$$f(z) = z^2 + \frac{1}{2}e^{(z-1)^2}, \quad a_1(z) = z^2 + \frac{1}{2} \quad \text{and} \quad a_2(z) = 3z - 1.$$

We note that

$$f(z) - \left(z^2 + \frac{1}{2}\right) = \frac{1}{2}(e^{(z-1)^2} - 1)$$

and

$$f'(z) - (3z - 1) = (z - 1)(e^{(z-1)^2} - 1).$$

Obviously  $f - a_1$  and  $f' - a_2$  share 0 IM, and  $\overline{N}_2(r, 0; f) \neq S(r, f)$  and  $\overline{N}(r, \infty; f) = 0$ , but  $f' \not\equiv a_2 a_1^{-1} f$ .

Example 1.7. Let

$$f(z) = \frac{2}{1 - e^{-2z}}.$$

Clearly  $f'(z) = -4e^{-2z}(1 - e^{-2z})^{-2}$ . We note that

$$f(z) - 1 = \frac{1 + e^{-2z}}{1 - e^{-2z}} \quad \text{and} \quad f'(z) - 1 = -\frac{(1 + e^{-2z})^2}{(1 - e^{-2z})^2}.$$

Obviously  $f$  and  $f'$  share 1 IM,  $\overline{N}(r, \infty; f) \neq S(r, f)$  and  $\overline{N}_2(r, 0; f) = 0$ , but  $f' \not\equiv f$ .

Example 1.8. Let  $f(z) = 1 + \tan z$ . Since  $\tan z$  does not assume the values  $\pm i$ , it follows that  $f(z)$  does not assume the values  $1 \pm i$ . So by the second fundamental theorem,  $\overline{N}(r, 0; f) = \overline{N}_2(r, 0; f) = T(r, f) + S(r, f)$  and  $\overline{N}(r, \infty; f) = T(r, f) + S(r, f)$ . Also we see that  $f'(z) - 1 = (f(z) - 1)^2$  and so  $f$  and  $f'$  share the value 1 IM, but  $f \not\equiv f'$ .

## 2. LEMMAS

In this section we present the lemmas which will be needed in the sequel.

**Lemma 2.1** ([4]). *Suppose that  $f$  is a transcendental meromorphic function and that*

$$f^n(z)P(f(z)) = Q(f(z)),$$

where  $P(f(z))$  and  $Q(f(z))$  are differential polynomials in  $f$  with functions of small proximity related to  $f$  as the coefficients and suppose that the degree of  $Q(f(z))$  is at most  $n$ . Then  $m(r, P) = S(r, f)$ .

**Lemma 2.2** ([7]). *Let  $f$  be a non-constant meromorphic function and let  $a_1(z)$ ,  $a_2(z)$  be two meromorphic functions such that  $T(r, a_i) = S(r, f)$ ,  $i = 1, 2$ . Then*

$$T(r, f) \leq \overline{N}(r, \infty; f) + \overline{N}(r, a_1; f) + \overline{N}(r, a_2; f) + S(r, f).$$

**Lemma 2.3** ([5]). *Let  $f(z)$  be a non-constant entire function and  $k (\geq 2)$  be an integer. If  $f(z)f^{(k)}(z) \neq 0$ , then  $f(z) = e^{az+b}$ , where  $a (\neq 0)$ ,  $b$  are constants.*

**Lemma 2.4.** *Let  $f$  be a non-constant meromorphic function such that*

$$(2.1) \quad (f^n)^{(k)} \equiv f^n,$$

where  $k, n \in \mathbb{N}$ . If  $n \geq k$ , then  $f$  assumes the form  $f(z) = ce^{z\lambda/n}$ , where  $c \in \mathbb{C} \setminus \{0\}$  and  $\lambda^k = 1$ .

*Proof.* First we suppose (2.1) holds. We claim that  $f$  does not have any pole. In fact, if  $z_0$  is a pole of  $f$  with multiplicity  $p$ , then  $z_0$  is a pole of  $f^n$  with multiplicity  $np$  and a pole of  $(f^n)^{(k)}$  with multiplicity  $np + k$ , which is impossible by (2.1). Hence  $f$  is a non-constant entire function. From (2.1), it is clear that  $f$  cannot be a polynomial. Therefore  $f$  is a transcendental entire function. We now consider the following two cases.

*Case 1.* Let  $n > k$ . If  $z_1$  is a zero of  $f$  with multiplicity  $q$ , then  $z_1$  is a zero of  $f^n$  with multiplicity  $nq$  and a zero of  $(f^n)^{(k)}$  with multiplicity  $nq - k$ , which is impossible by (2.1). Therefore from (2.1), we conclude that  $f^n(z)(f^n(z))^{(k)} \neq 0$ . If  $k \geq 2$ , then by Lemma 2.3 we have  $f(z) = ce^{z\lambda/n}$ , where  $c \in \mathbb{C} \setminus \{0\}$  and  $\lambda^k = 1$ .

Next we suppose  $k = 1$ . Since  $f(z) \neq 0, \infty$ , it follows that  $f(z) = e^{\alpha(z)}$ , where  $\alpha(z)$  is a non-constant entire function. Now from (2.1) we have  $\alpha'(z) = n^{-1}$ , i.e.,  $\alpha(z) = zn^{-1} + c_0$ , where  $c_0 \in \mathbb{C}$ . Consequently  $f(z) = ce^{z/n}$ , where  $c = e^{c_0}$ .

*Case 2.* Let  $n = k$ . First we suppose  $n = k = 1$ . Then from (2.1) we have  $f(z) \equiv f'(z)$  and so  $f(z) = ce^z$ , where  $c \in \mathbb{C} \setminus \{0\}$ .

Next we suppose  $n = k \geq 2$ . Let  $F = f^n$ . Then we have

$$\begin{aligned}
 (2.2) \quad F^{(k)} &= \frac{d^k}{dz^k}(f^k) = \frac{d^{k-1}}{dz^{k-1}}(kf^{k-1}f') = k \frac{d^{k-2}}{dz^{k-2}}((k-1)f^{k-2}(f')^2 + f^{k-1}f'') \\
 &= k(k-1) \frac{d^{k-2}}{dz^{k-2}}(f^{k-2}(f')^2) + k \frac{d^{k-2}}{dz^{k-2}}(f^{k-1}f'') \\
 &= k(k-1) \frac{d^{k-3}}{dz^{k-3}}((k-2)f^{k-3}(f')^3) + k(k-1) \frac{d^{k-3}}{dz^{k-3}}(2f^{k-2}f'f'') \\
 &\quad + k \frac{d^{k-3}}{dz^{k-3}}((k-1)f^{k-2}f'f'') + k \frac{d^{k-3}}{dz^{k-3}}(f^{k-1}f'') \\
 &= k(k-1)(k-2) \frac{d^{k-3}}{dz^{k-3}}(f^{k-3}(f')^3) + 2k(k-1) \frac{d^{k-3}}{dz^{k-3}}(f^{k-2}f'f'') \\
 &\quad + k(k-1) \frac{d^{k-3}}{dz^{k-3}}(f^{k-2}f'f'') + k \frac{d^{k-3}}{dz^{k-3}}(f^{k-1}f'') \\
 &= \dots = k!(f')^k + R(f),
 \end{aligned}$$

where  $R(f)$  is a differential polynomial in  $f$  such that each term of  $R(f)$  contains  $f^m$  for some  $m$  ( $1 \leq m \leq n-1$ ) as a factor.

From (2.1), we observe that  $f$  cannot have any multiple zero. Let  $z_2$  be a simple zero of  $f$ . Clearly  $z_2$  is a zero of  $F$  of multiplicity  $k$ . From (2.1), it is clear that  $z_2$  is also a zero of  $F^{(k)}$ . On the other hand  $z_2$  is a zero of  $R(f)$ . Now from (2.2), we observe that  $z_2$  is a zero of  $f'$ , which is impossible. Therefore  $f$  cannot have any simple zero. Hence  $f$  does not have any zero. Since from (2.1) we see that  $(f^n(z))^{(k)} f^n(z) \neq 0$ , by Lemma 2.3 we have  $f(z) = ce^{z\lambda/n}$ , where  $c \in \mathbb{C} \setminus \{0\}$  and  $\lambda^k = 1$ . This completes the proof.  $\square$

### 3. PROOFS OF THE THEOREMS

*Proof of Theorem 1.1.* Let

$$(3.1) \quad F = f^n.$$

Since  $S(r, f^n) = S(r, f)$ , from Lemma 2.2 we see that

$$nT(r, f) \leq \overline{N}(r, 0; F) + \overline{N}(r, a_1; F) + S(r, f^n) = \overline{N}(r, 0; f) + \overline{N}(r, a_1; F) + S(r, f).$$

Since  $n \geq k+1$ , it follows that  $\overline{N}(r, a_1; F) \neq S(r, f)$ . As  $F - a_1$  and  $F^{(k)} - a_2$  share “ $(0, 1)$ ”, it follows that  $\overline{N}(r, a_2; F^{(k)}) \neq S(r, f)$ .



Let  $z_0$  be a common zero of  $F - a_1$  and  $F^{(k)} - a_2$  such that  $a_i(z_0) \neq 0, \infty$  (otherwise the reduced counting functions of those zeros of  $F - a_1$  and  $F^{(k)} - a_2$  which are the zeros or poles of  $a_1(z)$  and  $a_2(z)$ , respectively, are equal to  $S(r, f)$ ), where  $i = 1, 2$ . Clearly  $F(z_0), F^{(k)}(z_0) \neq 0$ . Suppose  $z_0$  is a zero of  $F - a_1$  of multiplicity  $p_0$ . Since  $F - a_1$  and  $F^{(k)} - a_2$  share “(0, 1)”, it follows that  $z_0$  must be a zero of  $F^{(k)} - a_2$  of multiplicity  $q_0$ . Then in some neighbourhood of  $z_0$ , we get by Taylor’s expansion

$$\begin{aligned} F(z) &= a_{10} + a_{1r_0}(z - z_0)^{r_0} + a_{1r_0+1}(z - z_0)^{r_0+1} + \dots, \quad a_{10} \neq 0, \\ a_1(z) &= b_{10} + b_{1s_0}(z - z_0)^{s_0} + b_{1s_0+1}(z - z_0)^{s_0+1} + \dots, \quad b_{10} \neq 0. \end{aligned}$$

Since  $z_0$  is a zero of  $F - a_1$  of multiplicity  $p_0$ , it follows that  $a_{10} = b_{10}$  and  $p_0 \geq \min\{r_0, s_0\}$ , when  $p_0 = r_0 = s_0$ . Let us assume that

$$F(z) - a_1(z) = c_{1p_0}(z - z_0)^{p_0} + c_{1p_0+1}(z - z_0)^{p_0+1} + \dots, \quad c_{1p_0} \neq 0.$$

Therefore

$$\frac{F(z) - a_1(z)}{a_1(z)} = O((z - z_0)^{p_0}) \quad \text{and so} \quad \frac{F(z)}{a_1(z)} - 1 = O((z - z_0)^{p_0}).$$

Similarly

$$\frac{F^{(k)}(z) - a_2(z)}{a_2(z)} = O((z - z_0)^{q_0}) \quad \text{and} \quad \frac{F^{(k)}(z)}{a_2(z)} - 1 = O((z - z_0)^{q_0}).$$

Finally we conclude that  $F - a_1$  and  $F^{(k)} - a_2$  share “(0, 1)” if and only if  $F a_1^{-1}$  and  $F^{(k)} a_2^{-1}$  share “(1, 1)” except for the zeros and poles of  $a_1(z)$  and  $a_2(z)$ , respectively.

Let  $F_1 = f^n a_1^{-1}$  and  $G_1 = (f^n)^{(k)} a_2^{-1}$ . Clearly  $F_1$  and  $G_1$  share “(1, 1)” except for the zeros and poles of  $a_1(z)$  and  $a_2(z)$ , respectively, and so  $\overline{N}(r, 1; F_1) = \overline{N}(r, 1; G_1) + S(r, f)$ . Let

$$(3.2) \quad \Phi = \frac{F_1'(F_1 - G_1)}{F_1(F_1 - 1)} = \frac{F_1'}{F_1 - 1} \left(1 - \frac{G_1}{F_1}\right) = \frac{F_1'}{F_1 - 1} \left(1 - \frac{a_1}{a_2} \cdot \frac{F^{(k)}}{F}\right).$$

We now consider the following two cases.

*Case 1.* Let  $\Phi \not\equiv 0$ . Then clearly  $G_1 \not\equiv F_1$ , i.e.,  $(f^n)^{(k)} \not\equiv a_2 a_1^{-1} f^n$ . Now from (3.2) we get  $m(r, \infty; \Phi) = S(r, f)$ .

Let  $z_1$  be a zero of  $f$  of multiplicity  $p$  such that  $a_i(z_1) \neq 0, \infty$ , where  $i = 1, 2$ . Then  $z_1$  will be a zero of  $F_1$  and  $G_1$  of multiplicities  $np$  and  $np - k$ , respectively, and so from (3.2) we get

$$(3.3) \quad \Phi(z) = O((z - z_1)^{np-k-1}).$$

Since  $n \geq k + 1$ , it follows that  $\Phi$  is holomorphic at  $z_1$ .

Let  $z_2$  be a common zero of  $F_1 - 1$  and  $G_1 - 1$  such that  $a_i(z_2) \neq 0, \infty$ , where  $i = 1, 2$ . Suppose  $z_2$  is a zero of  $F_1 - 1$  of multiplicity  $q$ . Since  $F_1$  and  $G_1$  share “(1, 1)” except for the zeros and poles of  $a_1(z)$  and  $a_2(z)$ , respectively, it follows that  $z_2$  must be a zero of  $G_1 - 1$  of multiplicity  $r$ . Then in some neighbourhood of  $z_2$ , we get by Taylor’s expansion

$$\begin{aligned} F_1(z) - 1 &= b_q(z - z_2)^q + b_{q+1}(z - z_2)^{q+1} + \dots, \quad b_q \neq 0, \\ G_1(z) - 1 &= c_r(z - z_2)^r + c_{r+1}(z - z_2)^{r+1} + \dots, \quad c_r \neq 0. \end{aligned}$$

Clearly

$$F_1'(z) = qb_q(z - z_2)^{q-1} + (q + 1)b_{q+1}(z - z_2)^q + \dots$$

Note that

$$F_1(z) - G_1(z) = \begin{cases} b_q(z - z_2)^q + \dots & \text{if } q < r, \\ -c_r(z - z_2)^r - \dots & \text{if } q > r, \\ (b_q - c_q)(z - z_2)^q + \dots & \text{if } q = r. \end{cases}$$

Clearly from (3.2) we get

$$(3.4) \quad \Phi(z) = O((z - z_2)^{t-1}),$$

where  $t \geq \min\{q, r\}$ . Now from (3.4), it follows that  $\Phi$  is holomorphic at  $z_2$ .

We note from (3.2) that if  $z_*$  is a zero of  $F_1 - 1$  that is also a zero of  $a_2$  with multiplicity  $p_1$ , then  $z_*$  is a possible pole of  $\Phi$  with multiplicity at most  $1 + p_1$ . Again if  $z^*$  is a zero of  $f$  that is also a zero of  $a_2$  with multiplicity  $p_2$ , then  $z^*$  is a possible pole of  $\Phi$  with multiplicity at most  $k + p_2$ . So from (3.2), the above discussion and the hypothesis of Theorem 1.1 we note that

$$\begin{aligned} N(r, \infty; \Phi) &\leq (k + 1)N\left(r, \frac{a_1}{a_2}\right) + (k + 1)N(r, 0; a_1) + (k + 1)N(r, 0; a_2) \\ &\quad + (k + 1)\overline{N}(r, F_1) + (k + 1)\overline{N}(r, f) \\ &= (k + 1)\overline{N}(r, F_1) + S(r, f) = S(r, f). \end{aligned}$$

Consequently  $T(r, \Phi) = S(r, f)$ .

Let  $q \geq 2$ . Since  $F_1$  and  $G_1$  share “(1, 1)” except for the zeros and poles of  $a_1(z)$  and  $a_2(z)$ , it follows that  $r \geq 2$ . Therefore from (3.4) we see that

$$\overline{N}_{(2)}(r, 1; F_1) \leq N(r, 0; \Phi) + S(r, f) \leq T(r, \Phi) + S(r, f) = S(r, f).$$

Since  $F_1$  and  $G_1$  share “(1, 1)” except for the zeros and poles of  $a_1(z)$  and  $a_2(z)$ , it follows that  $\overline{N}_{(2)}(r, 1; G_1) = S(r, f)$ . Again from (3.2) we get

$$\frac{1}{F_1} = \frac{1}{\Phi} \left( \frac{F_1'}{F_1 - 1} - \frac{F_1'}{F_1} \right) \left( 1 - \frac{a_1}{a_2} \frac{(f^n)^{(k)}}{f^n} \right)$$

and so  $m(r, 1/F_1) = S(r, f)$ . Hence

$$(3.5) \quad m(r, 0; f) = m\left(r, \frac{1}{f}\right) = S(r, f).$$

We consider the following two sub-cases.

*Sub-case 1.1.* Let  $n > k + 1$ . From (3.3) we see that  $N(r, 0; f) \leq N(r, 0; \Phi) \leq T(r, \Phi) + O(1) = S(r, f)$ . Then from (3.5) we get  $T(r, f) = S(r, f)$ , which is a contradiction.

*Sub-case 1.2.* Let  $n = k + 1$ . Since for  $p \geq 2$ , we have  $np - k - 1 = (k + 1)p - k - 1 \geq p$ , from (3.3) we see that

$$N_{(2)}(r, 0; f) \leq N(r, 0; \Phi) \leq T(r, \Phi) + O(1) = S(r, f).$$

Then (3.5) gives

$$(3.6) \quad T(r, f) = N_{(1)}(r, 0; f) + S(r, f).$$

Note that  $\overline{N}_{(2)}(r, a_1; F) = \overline{N}_{(2)}(r, 1; F_1) + S(r, f) = S(r, f)$ ,  $\overline{N}_{(2)}(r, a_2; F^{(k)}) = \overline{N}_{(2)}(r, 1; G_1) + S(r, f) = S(r, f)$  and  $\overline{N}(r, \infty; F) = S(r, f)$ . Let

$$(3.7) \quad \beta = \frac{F^{(k)} - a_2}{F - a_1}, \quad \text{i.e., } F^{(k)} - a_2 = \beta(F - a_1).$$

We claim that  $\beta \neq 0$ . If not, suppose  $\beta \equiv 0$ . Then from (3.7) we have  $(f^n)^{(k)} \equiv a_2$ . Since  $n = k + 1$ , we immediately have  $N_{(1)}(r, 0, f) = S(r, f)$  and so from (3.6) we arrive at a contradiction. Hence  $\beta \neq 0$ . We now consider following two sub-cases.

*Sub-case 1.2.1.* Suppose  $T(r, \beta) \neq S(r, f)$ . Let  $z_{11}$  be a zero of  $F - a_1$  such that  $F^{(k)}(z_{11}) - a_2(z_{11}) \neq 0$ . Then obviously  $\beta$  has a pole at  $z_{11}$ . Let  $z_{12}$  be a zero of  $F^{(k)} - a_2$  such that  $F(z_{12}) - a_1(z_{12}) \neq 0$ . In that case  $\beta$  has a zero at  $z_{12}$ . Let  $z_{13}$  be a common zero of  $F - a_1$  and  $F^{(k)} - a_2$ . Since  $F - a_1$  and  $F^{(k)} - a_2$  share “(0, 1)”, it follows that  $\beta$  has a zero at  $z_{13}$  if  $z_{13}$  is a zero of  $F - a_1$  and  $F^{(k)} - a_2$  with multiplicities  $p_{13} (\geq 2)$  and  $q_{13} (\geq 2)$ , respectively, such that  $p_{13} < q_{13}$  and  $\beta$  has a pole at  $z_{13}$  if  $q_{13} < p_{13}$ . Therefore

$$\overline{N}(r, 0; \beta) \leq \overline{N}_{(2)}(r, a_2; F^{(k)}) + S(r, f) = S(r, f)$$

and

$$\overline{N}(r, \infty; \beta) \leq \overline{N}_{(2)}(r, a_1; F) + S(r, f) = S(r, f).$$

Let  $\xi = \beta'/\beta$ . Clearly

$$\begin{aligned} T(r, \xi) &= N\left(r, \infty; \frac{\beta'}{\beta}\right) + m\left(r, \frac{\beta'}{\beta}\right) \\ &= \overline{N}(r, 0; \beta) + \overline{N}(r, \infty; \beta) + S(r, \beta) = S(r, f) + S(r, \beta). \end{aligned}$$

Note that

$$\begin{aligned} T(r, \beta) &\leq T(r, F^{(k)} - a_2) + T(r, F - a_1) \leq T(r, F^{(k)}) + T(r, F) + S(r, F) + S(r, G) \\ &\leq (k+1)T(r, f^n) + nT(r, f) + S(r, f) = n(k+2)T(r, f) + S(r, f), \end{aligned}$$

which implies that  $S(r, \beta)$  can be replaced by  $S(r, f)$ . Consequently  $T(r, \xi) = S(r, f)$ . By logarithmic differentiation we get from (3.7)

$$(3.8) \quad \begin{aligned} F^{(k+1)}F - \xi F^{(k)}F - F^{(k)}F' &= a_1F^{(k+1)} - (\xi a_1 + a'_1)F^{(k)} - a_2F' \\ &\quad + (a'_2 - \xi a_2)F + \xi a_1 a_2 + a_2 a'_1 - a_1 a'_2. \end{aligned}$$

We deduce from (3.1) that

$$(3.9) \quad \begin{aligned} F^{(k)} &= \frac{d^k}{dz^k}(f^{k+1}) = \frac{d^{k-1}}{dz^{k-1}}((k+1)f^k f') = (k+1)\frac{d^{k-2}}{dz^{k-2}}(k f^{k-1}(f')^2 + f^k f'') \\ &= (k+1)k\frac{d^{k-2}}{dz^{k-2}}(f^{k-1}(f')^2) + (k+1)\frac{d^{k-2}}{dz^{k-2}}(f^k f'') \\ &= (k+1)k\frac{d^{k-3}}{dz^{k-3}}((k-1)f^{k-2}(f')^3) + (k+1)k\frac{d^{k-3}}{dz^{k-3}}(2f^{k-1}f' f'') \\ &\quad + (k+1)\frac{d^{k-3}}{dz^{k-3}}(k f^{k-1}f' f'') + (k+1)\frac{d^{k-3}}{dz^{k-3}}(f^k f''') \\ &= (k+1)k(k-1)\frac{d^{k-3}}{dz^{k-3}}(f^{k-2}(f')^3) + 2(k+1)k\frac{d^{k-3}}{dz^{k-3}}(f^{k-1}f' f'') \\ &\quad + (k+1)k\frac{d^{k-3}}{dz^{k-3}}(f^{k-1}f' f'') + (k+1)\frac{d^{k-3}}{dz^{k-3}}(f^k f''') \\ &= \dots = (k+1)!f(f')^k + \frac{k(k-1)}{4}(k+1)!f^2(f')^{k-2}f'' + \dots + (k+1)f^k f^{(k)}. \end{aligned}$$

Therefore

$$(3.10) \quad \begin{aligned} \frac{f'}{f}F^{(k)} &= (k+1)!(f')^{k+1} + \frac{k(k-1)}{4}(k+1)!f(f')^{k-1}f'' \\ &\quad + \dots + (k+1)f^{k-1}f'f^{(k)} \end{aligned}$$

and

$$(3.11) \quad \begin{aligned} F^{(k+1)} &= (k+1)!(f')^{k+1} + \frac{k(k+1)}{2}(k+1)!f(f')^{k-1}f'' \\ &\quad + \dots + (k+1)f^k f^{(k+1)}. \end{aligned}$$

Substituting (3.1), (3.9), (3.10) and (3.11) into (3.8), we have

$$(3.12) \quad f^n(z)P(z) = Q(z),$$

where  $Q(z)$  is a differential polynomial in  $f$  of degree  $n$  and

$$\begin{aligned}
 (3.13) \quad P(z) &= F^{(k+1)} - \xi F^{(k)} - n \frac{f'}{f} F^{(k)} \\
 &= -k(k+1)!(f')^{k+1} - (k+1)!\xi f(f')^k \\
 &\quad + \frac{k(k+1)(3-k)(k+1)!}{4} f(f')^{k-1} f'' \\
 &\quad + \dots + (k+1)f^k f^{(k+1)} - (k+1)\xi f^k f^{(k)} - (k+1)^2 f^{k-1} f' f^{(k)} \\
 &= -k(k+1)!(f')^{k+1} + R_1(f)
 \end{aligned}$$

is a differential polynomial in  $f$  of degree  $k+1$ , where  $R_1(f)$  is a differential polynomial in  $f$  such that each term of  $R_1(f)$  contains  $f^m$  for some  $m$  ( $1 \leq m \leq n-1$ ) as a factor.

We suppose that  $P \equiv 0$ . Then from (3.13) we get  $F^{(k+1)} - \xi F^{(k)} - nF^{(k)} f'/f \equiv 0$  and so  $F^{(k+1)}/F^{(k)} = \xi + n f'/f = \beta'/\beta + F'/F$ . By integration we have  $F^{(k)} = D\beta F$ , where  $D \in \mathbb{C} \setminus \{0\}$ . Since  $n = k+1$  and  $\overline{N}(r, \infty; \beta) = S(r, f)$ , it follows that  $\overline{N}(r, 0; f) = S(r, f)$ . Then from (3.6) we have  $T(r, f) = S(r, f)$ , which is a contradiction. So  $P \not\equiv 0$ . Then by Lemma 2.1 we get  $m(r, P) = S(r, f)$ . Since  $N(r, f) = S(r, f)$  we have

$$(3.14) \quad T(r, P) = S(r, f) \quad \text{and} \quad T(r, P') = S(r, f).$$

Note that from (3.13) we get

$$(3.15) \quad P'(z) = A_1(f')^k f'' + B_1(f')^{k+1} + S_1(f),$$

which is a differential polynomial in  $f$ , where  $A_1 = -\frac{1}{4}k(k+1)^2(k+1)!$ ,  $B_1 = -(k+1)!\xi$  and  $S_1(f)$  is a differential polynomial in  $f$  such that each term of  $S_1(f)$  contains  $f^m$  for some  $m$  ( $1 \leq m \leq n-1$ ) as a factor.

Let  $z_3$  be a simple zero of  $f$  such that  $\xi(z_3) \neq 0, \infty$ . Then from (3.13) and (3.15) we have

$$P(z_3) = -k(k+1)!(f'(z_3))^{k+1}, \quad P'(z_3) = A_1(f'(z_3))^k f''(z_3) + B_1(z_3)(f'(z_3))^{k+1}.$$

This shows that  $z_3$  is a zero of  $Pf'' - (K_1P' - K_2P)f'$ , where  $K_1 = -k(k+1)!/A_1$  and  $K_2 = B_1/A_1$ . Also  $T(r, K_1) = S(r, f)$  and  $T(r, K_2) = S(r, f)$ . Let

$$(3.16) \quad \Phi_1 = \frac{Pf'' - (K_1P' - K_2P)f'}{f}.$$

Then clearly  $m(r, \Phi_1) = S(r, f)$  and since  $N_{(2)}(r, 0; f) + N(r, f) = S(r, f)$ , we have  $T(r, \Phi_1) = S(r, f)$ . From (3.16) we obtain

$$(3.17) \quad f''(z) = \alpha_1(z)f(z) + \beta_1(z)f'(z),$$

where

$$(3.18) \quad \alpha_1 = \frac{\Phi_1}{P} \quad \text{and} \quad \beta_1 = K_1 \frac{P'}{P} - K_2.$$

Differentiating (3.17) and using it repeatedly we have

$$(3.19) \quad f^{(i)}(z) = \alpha_{i-1}(z)f(z) + \beta_{i-1}(z)f'(z),$$

where  $i \geq 2$  and  $T(r, \alpha_{i-1}) = S(r, f)$ ,  $T(r, \beta_{i-1}) = S(r, f)$ .

Also (3.18) yields

$$(3.20) \quad P' = \left( \frac{\beta_1}{K_1} + \frac{K_2}{K_1} \right) P \quad \text{and} \quad \beta_1 = K_1 \frac{P'}{P} - K_2 = \frac{-k(k+1)! P'}{A_1 P} - \frac{B_1}{A_1},$$

so that

$$(3.21) \quad A_1 \beta_1 + B_1 + k(k+1)! \frac{P'}{P} = 0.$$

Now we consider following two sub-cases.

*Sub-case 1.2.1.1.* Let  $k = 1$ . Now from (3.13) and (3.17) we have

$$P = -2(f')^2 - 2\xi f f' + 2f f'' = -2(f')^2 + (2\beta_1 - 2\xi) f f' + 2\alpha_1 f^2$$

and so

$$P' = (-2\beta_1 - 2\xi)(f')^2 + (2\beta_1' - 2\xi' + 2\beta_1^2 - 2\beta_1\xi) f f' + (2\alpha_1\beta_1 - 2\alpha_1\xi + 2\alpha_1') f^2.$$

Note that  $K_1 = 1$  and  $K_2 = \xi$  and so from (3.20) we have

$$(3.22) \quad (\beta_1' - \xi' - \beta_1\xi + \xi^2) f' + (-2\alpha_1\xi + \alpha_1') f \equiv 0.$$

If  $-2\alpha_1\xi + \alpha_1' \equiv 0$ , then from (3.22) we get, because  $f f' \not\equiv 0$ ,

$$(3.23) \quad \beta_1' - \xi' - \beta_1\xi + \xi^2 \equiv 0.$$

Let  $\beta_1 \equiv \xi$ . Then a simple calculation gives  $2\beta_1'\beta_1^{-1} = P'P^{-1}$  and so on integration we get  $\beta_1^2 = d_0P$ , where  $d_0 \in \mathbb{C} \setminus \{0\}$ . This contradicts the fact that  $T(r, \beta) \neq S(r, f)$ . So  $\beta_1 \not\equiv \xi$ . Now from (3.23) we get  $(\beta_1' - \xi')(\beta_1 - \xi)^{-1} = \xi = \beta_1'\beta_1^{-1}$ . So on integration we get  $\beta_1 = d_1(\beta_1 - \xi)$ , where  $d_1 \in \mathbb{C} \setminus \{0\}$ . This contradicts the fact that  $T(r, \beta) \neq S(r, f)$ . So we conclude that  $-2\alpha_1\xi + \alpha_1' \not\equiv 0$ . Then from (3.22) we see that if  $z_4$  is a simple zero of  $f$ , then  $z_4$  is either a pole of  $-2\alpha_1\xi + \alpha_1'$  or a zero of  $\beta_1' - \xi' - \beta_1\xi + \xi^2$ . Hence

$$N_1(r, 0; f) \leq N(r, \infty; -2\alpha_1\xi + \alpha_1') + N(r, 0; \beta_1' - \xi' - \beta_1\xi + \xi^2) = S(r, f).$$

So we arrive at a contradiction by (3.6).

*Sub-case 1.2.1.2.* Let  $k \geq 2$ . From (3.9) and (3.11) we have  $F^{(k)} = T_1(f)$ ,  $F^{(k+1)} = (k+1)!(f')^{k+1} + T_2(f)$  and  $F^{(k+2)} = \frac{1}{2}(k+1)(k+2)(k+1)!(f')^k f'' + T_3(f)$ , where  $T_1(f)$ ,  $T_2(f)$  and  $T_3(f)$  are differential polynomials in  $f$  such that each term of  $T_1(f)$ ,  $T_2(f)$  and  $T_3(f)$  contain  $f$  as a factor.

Comparing (3.8) and (3.12) and noting that  $F = f^n = f^{k+1}$  we have

$$(3.24) \quad \begin{aligned} Q &= a_1 F^{(k+1)} - (\xi a_1 + a'_1) F^{(k)} - a_2 F' + (a'_2 - \xi a_2) F + \gamma \\ &= a_1((k+1)!(f')^{k+1} + T_2(f)) - (\xi a_1 + a'_1) T_1(f) \\ &\quad - (k+1)a_2 f^k f' + (a'_2 - \xi a_2) f^{k+1} + \gamma, \end{aligned}$$

where  $\gamma = \xi a_1 a_2 + a_2 a'_1 - a_1 a'_2$ .

Now suppose  $\gamma(z) \equiv 0$ . Then by integration we obtain  $\beta = d_2 a_2 a_1^{-1}$ , where  $d_2 \in \mathbb{C} \setminus \{0\}$  and so  $T(r, \beta) = S(r, f)$ , which is a contradiction. Consequently  $\gamma(z) \not\equiv 0$ . Similarly we can verify that  $\xi a_1 + a'_1 \not\equiv 0$  and  $a'_2 - \xi a_2 \not\equiv 0$ . We further note that  $T(r, \gamma) = S(r, f)$ . Differentiating (3.24) we have

$$(3.25) \quad \begin{aligned} Q' &= a'_1 F^{(k+1)} + a_1 F^{(k+2)} - (\xi a_1 + a'_1) F^{(k+1)} - (\xi a_1 + a'_1)' F^{(k)} \\ &\quad - a'_2 F' - a_2 F'' + (a'_2 - \xi a_2)' F + (a'_2 - \xi a_2) F' + \gamma' \\ &= a'_1((k+1)!(f')^{k+1} + T_2(f)) - (\xi a_1 + a'_1)((k+1)!(f')^{k+1} + T_2(f)) \\ &\quad + a_1 \left( \frac{(k+1)(k+2)}{2} (k+1)!(f')^k f'' + T_3(f) \right) - (\xi a_1 + a'_1)' T_1(f) \\ &\quad - (k+1)a'_2 f^k f' - a_2(k(k+1)f^{k-1}(f')^2 + (k+1)f^k f'') \\ &\quad + (a'_2 - \xi a_2)' f^{k+1} + (k+1)(a'_2 - \xi a_2) f^k f' + \gamma'. \end{aligned}$$

Let  $z_5$  be a simple zero of  $f(z)$  such that  $z_5$  is not a zero or a pole of  $a_1$ ,  $a_2$  and  $\xi$ . Then from (3.12), (3.24) and (3.25) we have

$$\gamma(z_5) = A(z_5)(f'(z_5))^{k+1}, \quad \gamma'(z_5) = A_2(z_5)(f'(z_5))^k f''(z_5) + B_2(z_5)(f'(z_5))^{k+1},$$

where  $A(z) = -(k+1)!a_1(z)$ ,  $A_2(z) = -\frac{1}{2}(k+1)(k+2)(k+1)!a_1(z)$  and  $B_2(z) = (k+1)!\xi(z)a_1(z)$ . This shows that  $z_5$  is a zero of  $\gamma f'' - (K_3 \gamma' - K_4 \gamma) f'$ , where  $K_3 = AA_2^{-1}$  and  $K_4 = B_2 A_2^{-1}$ . Also  $T(r, K_3) = S(r, f)$  and  $T(r, K_4) = S(r, f)$ .

Let

$$(3.26) \quad \Phi_2 = \frac{\gamma f'' - (K_3 \gamma' - K_4 \gamma) f'}{f}.$$

Then clearly  $T(r, \Phi_2) = S(r, f)$ . From (3.26) we obtain

$$(3.27) \quad f'' = \varphi_1 f + \psi_1 f',$$

where

$$(3.28) \quad \varphi_1 = \frac{\Phi_2}{\gamma} \quad \text{and} \quad \psi_1 = K_3 \frac{\gamma'}{\gamma} - K_4.$$

Now we show that  $\psi_1 \neq \beta_1$ . If  $\psi_1 \equiv \beta_1$  then from (3.18) and (3.28) we have

$$\frac{2}{(k+1)(k+2)} \frac{\gamma'}{\gamma} + \frac{2}{(k+1)(k+2)} \xi \equiv \frac{4}{(k+1)^2} \frac{P'}{P} - \frac{4}{k(k+1)^2} \xi,$$

i.e.,

$$2k(k+2) \frac{P'}{P} - k(k+1) \frac{\gamma'}{\gamma} \equiv (k^2 + 3k + 4) \frac{\beta'}{\beta}.$$

On integration we have

$$\beta^{k^2+3k+4} \equiv \frac{d_3 P^{2k(k+2)}}{\gamma^{k(k+1)}},$$

where  $d_3 \in \mathbb{C} \setminus \{0\}$  and so from (3.14) we have  $T(r, \beta) = S(r, f)$ , a contradiction.

Now from (3.27) we have

$$(3.29) \quad f^{(i)} = \varphi_{i-1} f + \psi_{i-1} f',$$

where  $i \geq 2$  and  $T(r, \varphi_{i-1}) = S(r, f)$ ,  $T(r, \psi_{i-1}) = S(r, f)$ .

Also from (3.13), (3.15) and (3.29) we have, respectively,

$$(3.30) \quad P = -k(k+1)!(f')^{k+1} + \sum_{j=1}^{k+1} T_j f^j (f')^{k+1-j},$$

$$(3.31) \quad P' = (A_1 \psi_1 + B_1)(f')^{k+1} + \sum_{j=1}^{k+1} S_j f^j (f')^{k+1-j},$$

where  $T(r, T_j) = S(r, f)$  and  $T(r, S_j) = S(r, f)$ .

Multiplying (3.30) by  $P'$  and (3.31) by  $P$  and then subtracting we get

$$(3.32) \quad H_0 (f')^{k+1} + H_1 f (f')^k + \dots + H_{k+1} f^{k+1} \equiv 0,$$

where

$$(3.33) \quad H_0 = P \left( A_1 \psi_1 + B_1 + k(k+1)! \frac{P'}{P} \right) \quad \text{and} \quad H_j = P S_j - P' T_j \quad \text{for } j = 1, 2, \dots, k+1.$$

Since  $\beta_1 \neq \psi_1$  and  $P \neq 0$ , it follows from (3.21) and (3.33) that  $H_0 \neq 0$ . Again since  $H_0 (f')^{k+1} \neq 0$ , from (3.32) we conclude that  $H_i \neq 0$  for at least one  $i \in \{1, 2, \dots, k+1\}$ . Let  $S = \{1, 2, \dots, k+1\}$  and  $S_1 = \{i \in S: H_i \neq 0\}$ . Note that  $T(r, H_0) = S(r, f)$  and  $T(r, H_j) = S(r, f)$  for  $j \in S_1$ .

Now from (3.32) we see that a simple zero of  $f$  must be either a zero of  $H_0$  or a pole of at least one  $H_i$ 's, where  $i \in S_1$ . Therefore

$$N_{1\prime}(r, 0; f) \leq N(r, 0; H_0) + \sum_{j \in S_1} N(r, \infty; H_j) + S(r, f) = S(r, f).$$

So we arrive at a contradiction by (3.6).



*Sub-case 1.2.2.* Suppose  $T(r, \beta) = S(r, f)$ . Then from (3.7) we have

$$(3.34) \quad F^{(k)} - \beta F \equiv a_2 - \beta a_1.$$

If  $a_2 - \beta a_1 \equiv 0$ , then from (3.34) we get  $(f^n)^{(k)} \equiv a_2 a_1^{-1} f^n$ , which contradicts the fact that  $\Phi \not\equiv 0$ . So we suppose that  $a_2 - \beta a_1 \not\equiv 0$ . Let  $z_6$  be a simple zero of  $f$ . If  $z_6$  is not a pole of  $\beta$ , then from (3.34) we see that  $z_6$  is a zero of  $a_2 - a_1 \beta$ . Therefore

$$N_{11}(r, 0; f) \leq N(r, 0; a_2 - a_1 \beta) + N(r, \infty; \beta) = S(r, f).$$

So by (3.6) we arrive at a contradiction.

*Case 2.* Let  $\Phi \equiv 0$ . Now from (3.2) we get  $F_1 \equiv G_1$ , i.e.,  $(f^n)^{(k)} \equiv a_2 a_1^{-1} f^n$ .

Furthermore if  $a_1 \equiv a_2$ , then  $f^n \equiv (f^n)^{(k)}$ , and by Lemma 2.4,  $f$  assumes the form  $f(z) = ce^{z\lambda/n}$ , where  $c \in \mathbb{C} \setminus \{0\}$  and  $\lambda^k = 1$ .  $\square$

*Proof of Theorem 1.2.* Let  $F_1 = f^n a_1^{-1}$  and  $G_1 = (f^n)^{(k)} a_2^{-1}$ . Clearly  $F_1$  and  $G_1$  share “(1, 0)” except for the zeros and poles of  $a_1(z)$  and  $a_2(z)$  and so  $\overline{N}(r, 1; F_1) = \overline{N}(r, 1; G_1) + S(r, f)$ . We now consider following two cases.

*Case 1.* Let  $F_1 \not\equiv G_1$ . Then

$$(3.35) \quad \begin{aligned} \overline{N}(r, 1; F_1) &\leq \overline{N}(r, 0; G_1 - F_1 \mid F_1 \neq 0) + S(r, f) \\ &\leq \overline{N}\left(r, 0; \frac{G_1 - F_1}{F_1}\right) + S(r, f) \leq T\left(r, \frac{G_1 - F_1}{F_1}\right) + S(r, f) \\ &\leq T\left(r, \frac{G_1}{F_1}\right) + S(r, f) \leq N\left(r, \infty; \frac{G_1}{F_1}\right) + m\left(r, \infty; \frac{G_1}{F_1}\right) + S(r, f) \\ &= N\left(r, \infty; \frac{a_1 (f^n)^{(k)}}{a_2 f^n}\right) + m\left(r, \infty; \frac{a_1 (f^n)^{(k)}}{a_2 f^n}\right) + S(r, f) \\ &\leq k\overline{N}(r, \infty; f) + k\overline{N}(r, 0; f^n) + S(r, f) = k\overline{N}(r, 0; f) + S(r, f). \end{aligned}$$

Now using (3.35) and  $\overline{N}_{21}(r, 0; f) = S(r, f)$ , we get from the second fundamental theorem that

$$(3.36) \quad \begin{aligned} nT(r, f) &= T(r, f^n) + S(r, f) \leq T(r, F_1) + S(r, f) \\ &\leq \overline{N}(r, \infty; F_1) + \overline{N}(r, 0; F_1) + \overline{N}(r, 1; F_1) + S(r, F) \\ &\leq \overline{N}(r, \infty; f) + \overline{N}(r, 0; f^n) + \overline{N}(r, 1; F_1) + S(r, f) \\ &\leq (k+1)\overline{N}(r, 0; f) + S(r, f) \\ &\leq \frac{k+1}{3}N(r, 0; f) + S(r, f) \leq \frac{k+1}{3}T(r, f) + S(r, f). \end{aligned}$$

Since  $n \geq k$ , (3.36) leads to a contradiction.

*Case 2.*  $F_1 \equiv G_1$ . Then  $(f^n)^{(k)} \equiv a_2 a_1^{-1} f^n$ . Furthermore if  $a_1 \equiv a_2$ , then  $f^n \equiv (f^n)^{(k)}$ , and, by Lemma 2.4,  $f$  assumes the form  $f(z) = ce^{z\lambda/n}$ , where  $c \in \mathbb{C} \setminus \{0\}$  and  $\lambda^k = 1$ .  $\square$

## References

- [1] *R. Brück*: On entire functions which share one value CM with their first derivative. *Result. Math.* *30* (1996), 21–24. [zbl](#) [MR](#) [doi](#)
- [2] *T. Cao*: On the Brück conjecture. *Bull. Aust. Math. Soc.* *93* (2016), 248–259. [zbl](#) [MR](#) [doi](#)
- [3] *Z.-X. Chen, K. H. Shon*: On conjecture of R. Brück concerning the entire function sharing one value CM with its derivative. *Taiwanese J. Math.* *8* (2004), 235–244. [zbl](#) [MR](#) [doi](#)
- [4] *J. Clunie*: On integral and meromorphic functions. *J. Lond. Math. Soc.* *37* (1962), 17–27. [zbl](#) [MR](#) [doi](#)
- [5] *G. Frank*: Eine Vermutung von Hayman über Nullstellen meromorpher Funktionen. *Math. Z.* *149* (1976), 29–36. (In German.) [zbl](#) [MR](#) [doi](#)
- [6] *G. G. Gundersen, L.-Z. Yang*: Entire functions that share one value with one or two of their derivatives. *J. Math. Anal. Appl.* *223* (1998), 88–95. [zbl](#) [MR](#) [doi](#)
- [7] *W. K. Hayman*: *Meromorphic Functions*. Oxford Mathematical Monographs. Clarendon Press, Oxford, 1964. [zbl](#) [MR](#)
- [8] *I. Lahiri*: Weighted sharing and uniqueness of meromorphic functions. *Nagoya Math. J.* *161* (2001), 193–206. [zbl](#) [MR](#) [doi](#)
- [9] *I. Lahiri*: Weighted value sharing and uniqueness of meromorphic functions. *Complex Variables, Theory Appl.* *46* (2001), 241–253. [zbl](#) [MR](#) [doi](#)
- [10] *S. Lin, W. Lin*: Uniqueness of meromorphic functions concerning weakly weighted-sharing. *Kodai Math. J.* *29* (2006), 269–280. [zbl](#) [MR](#) [doi](#)
- [11] *F. Lü, H. Yi*: The Brück conjecture and entire functions sharing polynomials with their  $k$ -th derivatives. *J. Korean Math. Soc.* *48* (2011), 499–512. [zbl](#) [MR](#) [doi](#)
- [12] *W. Lü, Q. Li, C. Yang*: On the transcendental entire solutions of a class of differential equations. *Bull. Korean Math. Soc.* *51* (2014), 1281–1289. [zbl](#) [MR](#) [doi](#)
- [13] *S. Majumder*: A result on a conjecture of W. Lü, Q. Li and C. Yang. *Bull. Korean Math. Soc.* *53* (2016), 411–421. [zbl](#) [MR](#) [doi](#)
- [14] *C.-C. Yang, H.-X. Yi*: *Uniqueness Theory of Meromorphic Functions. Mathematics and Its Applications (Dordrecht)* 557. Kluwer Academic Publishers, Dordrecht, 2003. [zbl](#) [MR](#) [doi](#)
- [15] *L.-Z. Yang, J.-L. Zhang*: Non-existence of meromorphic solutions of Fermat type functional equation. *Aequationes Math.* *76* (2008), 140–150. [zbl](#) [MR](#) [doi](#)
- [16] *J. Zhang*: Meromorphic functions sharing a small function with their derivatives. *Kyungpook Math. J.* *49* (2009), 143–154. [zbl](#) [MR](#) [doi](#)
- [17] *J. Zhang, L. Yang*: A power of a meromorphic function sharing a small function with its derivative. *Ann. Acad. Sci. Fenn., Math.* *34* (2009), 249–260. [zbl](#) [MR](#)
- [18] *J.-L. Zhang, L.-Z. Yang*: A power of an entire function sharing one value with its derivative. *Comput. Math. Appl.* *60* (2010), 2153–2160. [zbl](#) [MR](#) [doi](#)

*Authors' addresses:* *Indrajit Lahiri*, Department of Mathematics, University of Kalyani, Kalyani, West Bengal 741235, India, e-mail: [ilahiri@hotmail.com](mailto:ilahiri@hotmail.com); *Sujoy Majumder*, Department of Mathematics, Raiganj University, Raiganj, West Bengal 733134, India, e-mail: [sm05math@gmail.com](mailto:sm05math@gmail.com), [sujoy.katwa@gmail.com](mailto:sujoy.katwa@gmail.com).